

Supplementary data

Collection, characterization and evaluation of wild *Hedysarum coronarium* L. populations from Andalusia (southern Spain)

Eva María Córdoba, Salvador Nadal, Belén Román, Clara Isabel González-Verdejo

Additional file 1. The frequency growth habits (%) of different sulla accessions.

Accessions code	Plant growth habit		
	Rosette	Prostrate	Erect
<i>Hc-1</i>	0	40	60
<i>Hc-3</i>	20	60	20
<i>Hc-4</i>	60	20	20
<i>Hc-5</i>	20	60	20
<i>Hc-6</i>	20	80	0
<i>Hc-7</i>	80	20	0
<i>Hc-8</i>	40	40	20
<i>Hc-9</i>	20	20	60
<i>Hc-10</i>	40	60	0
<i>Hc-12</i>	100	0	0
<i>Hc-13</i>	0	100	0
<i>Hc-14</i>	0	100	0
<i>Hc-15</i>	100	0	0
<i>Hc-16</i>	20	60	20
<i>Hc-17</i>	40	40	20
<i>Hc-18</i>	80	20	0
<i>Hc-23</i>	80	0	20
<i>Hc-24</i>	0	80	20
<i>Hc-25</i>	0	100	0
<i>Hc-27</i>	0	100	0
<i>Hc-29</i>	100	0	0
<i>Hc-30</i>	0	20	80
<i>Hc-31</i>	80	20	0
<i>Hc-32</i>	100	0	0
<i>Hc-33</i>	20	80	0
<i>Hc-34</i>	0	40	60
<i>Carmen</i> cv.	0	0	100
<i>Grimaldi</i> cv.	0	80	20
<i>Del País</i> cv.	0	60	40
<i>Average</i>		35,1	44,8
			20