

Participatory assessment of potato farming systems, production constraints and cultivar preferences in Uganda

Prossy Namugga*, Rob Melis Julia Sibiya, And Alex Barekye




Figure1: Location of the surveyed potato growing areas in Uganda

Supplementary Table 1: Importance of potato (% of respondents) in the eight districts

Region	District	Cash	Food	Both
South Western	Kabale	0.0	1.2	98.8
	Kisoro	0.9	1.8	97.3
	Kanungu	0.0	1.4	98.6
Central	Rakai	4.8	4.8	90.5
	Lwengo	8.9	4.5	86.6
Eastern	Kween	0.0	0.0	100.0
	Kapchorwa	0.0	2.4	97.6
	Mbale	0.0	3.6	96.4
Mean		4.9	3.1	93.3

Supplementary Table 2: Commonly grown potato varieties as ranked by farmers (% of respondents)

	Kabale	Kisoro	Kanungu	Rakai	Lwengo	Kween	Kapchorwa	Mbale
Rwangume	93.3	25.0	98.0	98.4	95.6	33.3	14.3	82.1
Victoria	25.0	6.3	12.9	14.3	2.9	31.3	14.8	50.0
Kinigi	29.9	87.5	34.3	0.0	0.0	0.0	0.0	0.0
Rutuku	1.8	0.0	5.7	0.0	0.0	0.0	0.0	0.0
Rwashaki	9.2	16.1	1.4	0.0	0.0	0.0	0.0	0.0
Kabera	0.0	0.0	0.0	0.0	2.9	0.0	0.0	0.0
Wanale	0.0	0.0	0.0	0.0	0.0	13.3	2.4	7.1
Kasumali	0.0	0.0	0.0	7.9	0.0	0.0	0.0	0.0
Sutama	0.0	16.1	0.0	0.0	0.0	0.0	0.0	0.0
Kimuri	5.5	0.0	1.4	0.0	0.0	0.0	0.0	0.0
Petero	0.0	6.3	0.0	0.0	0.0	0.0	0.0	0.0
Nakpot5	0.0	0.0	0.0	0.0	0.0	13.3	7.1	0.0
Cruza	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.6
Kabale	0.0	0.0	0.0	3.2	0.0	66.7	97.6	0.0
Singo	0.0	0.0	0.0	1.6	4.4	0.0	0.0	0.0
Agriculture	0.0	0.0	0.0	0.0	1.5	0.0	2.4	0.0
Makerere	0.0	0.0	0.0	0.0	7.4	0.0	0.0	0.0

Supplementary Table 3: Decision on when to spray (% of respondents in each district)

Decision	Kabale	Kisoro	Kanungu	Rakai	Lwengo	Kween	Kapchorwa	Mbale
After germination	61.6	81.6	63.7	2.4	19.4	30.6	16.4	12.5
Before 1st weeding	0.0	0.0	0.0	0.0	1.0	0.0	5.5	0.0
After weeding	13.2	13.2	5.5	42.9	25.5	41.7	36.4	46.9
At flowering	7.4	1.8	9.9	29.8	19.4	5.6	14.5	6.3
After flowering	5.8	0.9	13.2	19.0	20.4	2.8	7.3	3.1
When it is humid	1.6	0.0	6.6	1.2	4.1	0.0	0.0	3.1
NA	5.3	1.8	1.1	4.8	9.2	5.6	7.3	6.3
When symptoms appear	5.3	0.9	0.0	0.0	1.0	13.9	12.7	21.9