

## Comparative impact of Arbuscular Mycorrhizal fungi from Moroccan soils and commercial inoculum on potato yield and nutrient composition

Wissame CHAFAI\*, Halima Bouchentouf, Ahmed KHALID

Laboratory for the Improvement of Agricultural Production, Biotechnology and Environment, Faculty of Sciences, Mohammed Premier University, Oujda, Morocco

\*Corresponding author: [wissamesalma@gmail.com](mailto:wissamesalma@gmail.com)

### Abstract

Potato (*Solanum tuberosum* L.) is a widely grown crop in Morocco. Mycorrhiza fertilization of potatoes needs to be monitored in order to observe their effects on the growth of potatoes. The aim of this study was therefore to test the effect of fungal strains on potato yield and tuber quality. The inoculation treatments included a commercial inoculum (T2), a natural inoculum (T3), represented by leek roots fragments, obtained after trapping by this host plant of AMF spores naturally present in the soil of the region of Guercif-Morocco, and a control treatment consisting of a sterile substrate. Our results showed significant differences in the frequency and intensity of mycorrhization between the T3 inoculum and the T2 inoculum, while control plants showed no AMF infection. Compared to the T1 control (non-inoculated plants), the gains in number of tubers/plant was 24% and 39% higher, whereas gains in yield was 34% and 48% higher for potato plants inoculated with T2 and T3 inocula, respectively. Furthermore, chemical analyses showed the important role of controlled mycorrhization in the accumulation of mineral elements; N, P, K, Zn, Cu and Fe in potato tubers. The T3 inoculum obtained from the soil of Guercif-Morocco allowed to obtain the best value of dry matter (16.82 g/100 g fM) and the best contents in Zn; K and P and the tubers.

**Keywords:** Potato; endomycorrhiza; mineral composition of tubers.

**Abbreviations:** AMF\_Arbuscular mycorrhizal fungi; fM\_Fresh matter; N\_Nitrogen; K\_Potassium; P\_Phosphorus; Zn\_Zinc; Cu\_Copper; Fe\_Iron; Ca\_Calcium.

### Introduction

Potato (*Solanum tuberosum* L., Solanales Solanaceae) is the most important tuber in the world. The potato crop is largely grown due to its valuable marketing (AL-Shmary and AL-Taey, 2020). It is widely grown in more than 125 countries and consumed daily by more than one billion people (FAO, 2015). Potato (*Solanum tuberosum*) occupies an important place in the staple diet of people in developing countries. Since its introduction to Morocco in the 19th century, the potato has become an increasingly important part of the Moroccan diet. Its cultivation occupies an area of up to 60,000 hectare (ha) per year, equivalent to 23% of the total market garden area resulting in a production of 1.5 million tons annually. From a nutritional point of view, it is one of the most nutritious tuber plants with a high energy content (Hassan, 2003). However, this crop does not always yield as expected since it is one of the most demanding crops in terms of chemical fertilizer inputs compared to other vegetable crops. Its nitrogen (N), phosphate (P) and potassium (K) requirements are 100%, 100% and 33% higher than other vegetable crops, respectively (Khoshnevisan et al., 2013). However, these chemical fertilizers have some adverse effects on the environment and human health (Kumar and Keshar, 2017). For this reason, it is necessary to find safer alternatives to these chemical fertilizers to improve soil fertility and plant growth.

According to many researchers, a shift towards a more productive agriculture that is less dependent on chemical

phosphate inputs cannot be achieved without better management of biological interactions in agrosystems (Plenchette et al., 2005). Mycorrhizae, symbiotic associations between soil fungi and plant roots, are an example of such agrosystems. These fungi form symbiotic associations with almost 80% of plant species, including the majority of staple crops (Cobb et al. 2021; Thirkell et al. 2022).

The beneficial effects of arbuscular mycorrhizal symbiosis on plant growth and health are well documented (Begum et al., 2019). It increases their resistance and/or tolerance to numerous abiotic and biotic stresses (Sánchez-Romera et al., 2016; Ravnskov et al., 2020). In fact, arbuscular mycorrhizal fungi (AMF) have been shown to improve soil fertility, promote increased phosphorus (P) uptake by plants (Püschel et al., 2021), as well as the uptake of other essential nutrients such as N, S, Cu and Zn (Bhandari and Garg 2017; Ruytinx et al. 2020).

The use of AMF has been the subject of previous studies (Wu et al., 2013; Etlili et al., 2022; Bousageon et al., 2023) that showed the effectiveness of these microorganisms on improvement of potato growth.

Although mycorrhizal fungi are ubiquitous in natural soils, cultural practices most often disrupt their populations and diversity (Jansa et al., 2003). Under these conditions, numerous experiments conducted for decades have shown that the addition of spores or propagules of these fungi to the soil results in faster plant development and increased yields.

It is in this context that the present study was initiated and its objective is to evaluate the effectiveness, in the short term, of a commercial product/inoculum that is composed by the genus *Glomus*, and of a natural inoculum constituted by fragments of leek roots having trapped AMF spores naturally present in the soil of the region of Guercif-Morocco on the growth, the yield and the nutritious quality of potato.

## Results

### **Mycorrhizal colonization of plant roots**

Compared to the control (non-inoculated plants; T1), the mycorrhization rates of potato was 100% and 78% for the natural inoculum from Guercif-Maroc soil and the commercial inoculum, respectively. No presence of mycorrhizal fungi was observed in the roots of non-inoculated control plants.

The impact of the application of the different mycorrhizal inocula was reflected in different frequencies of mycorrhizal colonization (48% for plants inoculated with the commercial inoculum (T2) and 84% for plants inoculated with naturel inoculum (T3)). The best frequency was recorded for the T3 inoculum and was significantly different from the T2 treatment (Figure 1).

The colonization intensity of potato plant roots was significantly higher in T2 (NI) than T3 (CI) (52% in T2 vs. 24% in T3) (Figure 1). These results suggest the presence of more active AMF species in the inoculum from the Guercif-Morocco soil (T2) than in the commercial inoculum (T3).

### **Richness and diversity of AMF**

The density of AMF spores extracted from the potato culture substrate inoculated with T3 inoculum at the end of the experiment was lower (201spores/100g substrate) than that found in the Guercif -Morocco soil (3860/100g soil) that was used for the preparation of the T3 inoculum (Chafai et al.,2022).

The morphological study of the spores revealed the presence of seven different morphotypes represented by the following species: *Funneliformis constructum*; *Rhizophagus clarus*; *Clomus macrocarpum*; *Rhizophagus intaradices*; *Glomus deserticola*; *Funneliformis mosseae* and *Claroideoglomus etunicatum* (Table 1). These species belonged to the family Glomaceae, and the order Glomerales.


*Rhizophagus intraradices* and *Funneliformis mosseae* were the most frequent species from the T3 inoculated culture substrate (Figure 3).

### **Effect of mycorrhizae on yield**

Compared to non-inoculated control plants, Mycorrhizal inoculation resulted in an increase of the potato's total yield, which was 34% higher with the T2 inoculum and 48% higher with the T3 inoculum.

The best yield (244g tubers/plant) was recorded in plants inoculated with natural inoculum isolated from the Guercif-Morocco soil (T3). These results show a significant difference between the two inocula (T1 and T3) (Figure 4).

The number of tubers per plant were improved by mycorrhization (Figure 5). T3 significantly increased the number of tubers while T2 were not significantly higher than the control. Plants mycorrhized by T2 and T3 increased their number of tubers per plant by about 24% and 39%, respectively, compared to non-mycorrhized control plants (T1).



**Figure 1.** Frequency and intensity of mycorrhization of potato roots. (T1: plant control; T2: Plants inoculated with commercial inoculum; T3: Plants inoculated with natural inoculum). Each of the mycorrhization parameters (frequency and intensity) was treated statistically independently of the other. Treatments sharing the same letter are not significantly different ( $p < 0.05$ ).

### **Tuber nutrient uptake**

Inoculation with commercial inoculum (T2) and natural inoculum (T3) significantly increased the potato tuber biomass by approximately 2.3% and 2.9% for T2 and T3, respectively (Table 2).

Mycorrhization significantly improved the N content of the tubers. In T2, The N content of the tubers was almost 10 times that of the control.

The results presented in Table 2 show that inoculation of potato plants with the different inocula (T2 and T2) had a positive influence on the content of Ca; Zn; Cu; P, Fe and K in the tubers.

Compared to the non-treated control, treatment with T2 resulted in a significant increase of the Ca content of the potato tubers, while the inoculation with T3 resulted in a significant decrease in the Ca content.

The highest Zn content was recorded for T3 (0.76%) with a significant difference from T2 (0.586%) and T1 (0.497%) (Table 2).

The highest Cu content was found for T2 followed by T3 and T1 with a significant difference.

T2 and T3 had higher Fe content than the control (T1) with a significant statistical difference.


The highest K content (77.199%) was recorded for T3 compared to T2 and T1 (71.934% and 69.221 respectively). The inoculated plants showed a significant difference from the non-inoculated plants (T1) (Table 3).

Mycorrhization increased the P content of the inoculated plants compared to the control plants where the highest value was observed in the T3 treatment.

The S and Si contents are higher in the control than in the inoculated plants.

## Discussion

While the roots of the potato plants were well mycorrhized by both the commercial inoculum T2 and the natural inoculum T3 (isolated from the Guercif -Morocco soil), No mycorrhization was observed in the non-inoculated T1 control plants in our experimental set-up, indicating the absence of endogenous fungi in our growing medium. The contribution of the mycorrhizal inocula resulted in the formation of different structures common to endomycorrhizae (Béreau and Garbaye, 1994).



**Figure 2.** Structures of endomycorrhizal fungi observed in root samples of potato plants inoculated with commercial inoculum (T2) and natural inoculum (T3) and non-inoculated plants (T1): arbuscules (a), vesicles (v), spores (s), and extra and intracellular hyphae (he), (hi) (G×400).

**Table 1.** Characteristics of all AMF isolated from the culture substrate inoculated with T3 of inoculum.


Species	form	Color	Size of the spore (µm)	Wall size (µm)	Length of the hypha (µm)	Surface of the spore
<i>Funneliformis constructum</i>	globular	Almost black	110	1	-	smooth
<i>Rhizophagus Clarus</i>	globular	Brown	96	1		smooth
<i>Clomus macrocarpum</i>	globular	Yellow-brown	104	1.1	-	smooth
<i>Rhizophagus intraradices</i>	globular	Hyaline	96	1.5	99	smooth
<i>Glomus deserticola</i>	globular	Reddish brown	40	1.3	-	smooth
<i>Funneliformis mosseae</i>	globular	Yellow	90	1.2	-	smooth
<i>Claroideoglomus etunicatum</i>	globular	Orange	100	2	-	smooth

The addition of AMF inoculum to the substrate positively affected the yield parameters of the potato crop. Indeed, the number of tubers and yield of inoculated plants were much higher than those of non-inoculated plants. In fact, we showed that the number of tubers/potato plant increased by 24% and 39% for plants inoculated with T2 and T3 respectively. In addition, the yield of plants inoculated with T2 and T3 were increased by 34% and 48% respectively. Our results corroborate that a recent study by Lombardo et al. (2020), who showed an increase in marketable tuber yield by 25% and in tuber/plant number by 21% for inoculated plants. Tohidi-Moghaddam et al. (2004) reported that by solubilizing phosphates, the microorganisms would increase available phosphorus in the soil, which could increase the number of tubers per plant. This suggests that the parameters for estimating the degree of root colonization vary among treatments, and optimization of symbiotic interaction may depend on the selection of appropriate AMF isolates for specific cultivars. In fact, the mycorrhization frequency (F) and mycorrhizal intensity (M) values obtained with the T3 inoculum (84% and 52%, respectively), were higher than those obtained with the T2 inoculum (48% and 24%, respectively). This suggests that our inoculum isolated from Guercif-Morocco soil (T3) has a higher mycorrhizogenic potential when compared to commercial inoculum (T2). These interesting results obtained with the T3 inoculum could be the result of the development of a larger network of hyphae in the rhizosphere of inoculated plants, which was


also shown in previous studies (Zhong-Qun et al. (2007) and Tahat et al. (2008).

The assessment of the biodiversity of mycorrhizal fungi in the rhizosphere of potato plants that received T3 inoculum showed the presence of seven morphotypes. The majority of these species belong to the genus *Glomus* which generally dominates AMF communities in Moroccan agricultural soils (Sghir et al., 2014; Chliyeh et al., 2014; Talbi et al., 2015; Selmaoui et al., 2017). The other species identified in the Guercif-Maroc soil, which was the source of the T3 inoculum (Chafai et al., 2022) have disappeared, or have stopped their activity. Several factors can affect the development of AMF communities in the soil. For example, early colonization of host plant roots by one AMF species may prevent further colonization by other fungi. This phenomenon is known as self-regulation (Hepper et al., 1988; Person et al., 1993; Vierheilig et al., 2000; Vierheilig, 2004).


*Rhizophagus intraradices* and *Funneliformis mosseae* are the most frequent AMF fungal species in the substrate of our potato crop. A similar result was obtained by Bharadwaj et al. (2007), who showed, that when using a potato trap crop, *Rhizophagus intraradices* and *Funneliformis mosseae* are the most common fungi in the host plant roots and the crop soil. The diversity of AMF species in the potato growing medium that received the T3 inoculum, was found to be effective on the growth and yield of the host plant under our growing conditions. Ortas and Ustuner (2014), also showed that the richness and interaction of AMF species positively impacted plant growth. Functional complementarity between species



**Figure 3.** Frequency of occurrence of AMF species in potato crop substrate inoculated with natural inoculum (T3) at the end of the experiment. Treatments sharing the same letter are not significantly different ( $p < 0.05$ ).



**Figure 4:** Potato crop yields. T1: Control-non-mycorrhized plants; T2: Plants inoculated with commercial inoculum; T3: Plants inoculated with natural inoculum. Treatments sharing the same letter are not significantly different ( $p < 0.05$ ).



**Figure 5.** Number of potato tubers (T1: Control; T2: Plants inoculated with commercial inoculum; T3: Plants inoculated with natural inoculum). Treatments sharing the same letter are not significantly different ( $p < 0.05$ ).

in the AM community colonizing a single root system has been suggested by Jansa et al., 2008; Koide, 2000. Similar to previous studies by Vosatka and Gryndker (2000) and Ndiaye et al. (2005), our study showed that potato plants mycorrhized by T2 and T3 had a higher dry biomass production compared to the non-mycorrhized control. The addition of AMF T2 and T3 to the growing medium, significantly modified the content of potato tubers for many mineral elements such as Zn; Cu; Fe; P, N and K. As already shown by different previous studies that highlight the

importance of AMF in improving the absorption of mineral elements (nitrogen, phosphorus, potassium and microelements) with low mobility in the soil by tomato, (Hart et al.2015; Sánchez-Bel et al.2018; Higo et al.2020). Nitrogen was the mineral element with the highest content in inoculated plants compared to uninoculated plants, while S and Si were not affected by mycorrhization and the values recorded in uninoculated plants exceeded those in mycorrhized plants. Our work clearly showed that the T3 inoculum resulted in an increase in P, K, Zn and Fe content of potato tubers compared to the T2 commercial inoculum. Our results confirmed the important role of adding AMF to the crop soil. This makes these microorganisms an eco-sustainable tool to increase yield and increase the levels of macro and micro mineral elements in potato tubers.

## Materials and methods

### Producing natural inoculum

A composite endomycorrhizal inoculum was prepared from the soil of Guercif-Maroc (33°58'45"N, 3°15'41"W, Altitude: 688 m). AMF trapping was performed with leek plants as host plants. After 16 weeks of culture, the roots of the seedlings were excised, rinsed with distilled water and cut into 1-2 mm fragments. These root fragments were used as endomycorrhizal inoculum (T3). The inoculum consisted of 31 endomycorrhizal fungal species, identified morphologically by Chafai et al (2022).

### Plant cultivation and treatment

A pot experiment with potato was carried out under semi-controlled conditions in order to evaluate the effectiveness of inoculation of potato plants with two types of endomycorrhizal fungi inocula and to compare the effects of these inocula with those of non-inoculated control plants. Potato seed tubers were sterilized with 10% H<sub>2</sub>O<sub>2</sub> for 10 min to remove surface fungi, and then rinsed thoroughly with water.

The inoculation was carried out at the time of sowing, on sterilized substrate, according to the following experimental protocol:

- T1: non-inoculated control plants.
- T2: Plants inoculated with a commercial liquid multi-species inoculum (genus Glomus).
- T3: The plants inoculated with a natural inoculum, in the form of fragments of mycorrhized leek roots, obtained by trapping AMF spores naturally present in the soil of the Guercif-Morocco region.

Each treatment had three replicates, which were placed in a completely randomized design.

No fertilizers were applied during the entire growing cycle. However, three treatments against downy mildew were applied.

### Experimental materials

The variety of potato used is "Spunta". The substrate used for the pot experiment was a mixture of sand and peat (v/v) and it was autoclaved for 20 min at 120 °C. Sand was considered an inert substrate and a peat had a super fine structure; pH 6.1; salinity 1.2 g L<sup>-1</sup>; N 210 mg L<sup>-1</sup>; P,O 120 mg L<sup>-1</sup>; K,O 260 mg L<sup>-1</sup>; NPK balance 1-0.6-1.2.

The cylindrique pots used for the experiment had a volume of 7L with 20 cm inner diameter and 24 cm in height. After pouring the substrate into each pot, the height of the soil was 20 cm.

**Table 2.** Potato tuber dry matter (g/100gMf), nitrogen and mineral element content (%/ g dry matter).

Traitement	dry matter (g/100g FM)	N (mg/gMs)	Mineral element in %							
			Ca%	Zn%	Cu%	Fe%	K%	S%	P%	Si%
T1	16.34 <sup>b</sup>	26,75 <sup>c</sup>	13.958 <sup>b</sup>	0.497 <sup>c</sup>	0.922 <sup>c</sup>	1.069 <sup>b</sup>	69.221 <sup>c</sup>	6.785 <sup>a</sup>	2.516 <sup>b</sup>	4.972 <sup>a</sup>
T2	16.72 <sup>a</sup>	216.87 <sup>a</sup>	14.517 <sup>a</sup>	0.586 <sup>b</sup>	1.39 <sup>a</sup>	1.731 <sup>a</sup>	71.934 <sup>b</sup>	5.053 <sup>b</sup>	2.676 <sup>ab</sup>	2.111 <sup>b</sup>
T3	16.82 <sup>a</sup>	117.5 <sup>b</sup>	9.971 <sup>c</sup>	0.761 <sup>a</sup>	1.155 <sup>b</sup>	1.759 <sup>a</sup>	77.199 <sup>a</sup>	4.38 <sup>b</sup>	3.013 <sup>a</sup>	1.761 <sup>b</sup>

T1: Non-inoculated plants; T2: plants inoculated with commercial inoculum; T3: Plants inoculated with natural inoculum. Two values in the same column followed by the same letter are not significantly different at the 5% level.

### Evaluation of root colonization

Potato root samples were taken at the end of the crop and the assessment of mycorrhization parameters was performed by observing randomly selected root fragments of about 1 cm (Trouvelot and kough, 1986; Amir and renard, 2003). The frequency (F %) and mycorrhizal intensity (M%) of root MACs were measured by assigning a mycorrhizal index ranging from 0 to 5 (Derkowska et al., 2008).

- Mycorrhization frequency (F): extent of infection of the root system

$$F\% = \left( \frac{N - N_0}{N} \right) * 100$$

N = number of observed fragments and N<sub>0</sub> = number of fragments without any trace of mycorrhization.

- Colonization intensity (M): It expresses the portion of the colonized cortex compared to the whole root system.

$$M\% = [(95 * n_5) + (70 * n_4) + (30 * n_3) + (5 * n_2 + n_1)] / N$$

N: number of observed fragments

Where n<sub>5</sub>, n<sub>4</sub>,...n<sub>1</sub> are the number of fragments noted 5, 4,..., 1 respectively; class 5: more than 91%, class 4: from 51 to 90%, class 3: from 11 to 50%, class 2: less than 10%, class 1: trace and class 0: no mycorrhization.

### Extraction and identification of spores from the culture medium (T3)

Extraction of spores of endomycorrhizal fungi from the substrate was performed according to the method of Gerdemann and Nicolson (1963). The number of spores in 100 g of substrate was estimated by direct counting under a binocular loupe. Spores were then observed between slide and coverslip under a light microscope (x40) and classified according to color, shape, wall structure, and hyphal attachments to identify their genus (Ferrer and Herrera 1981; Schenck and Smith 1982; Walker and Mize 1982; Hall 1984; Schenck and Perez 1987; Morton and Benny 1990; and information available in various databases (INVAM 2017)).

Isolated spores were quantified by direct counting under a binocular loupe to estimate the number in 100 g of substrate (spore densities). Three replicates were performed.

The frequency of appearance of the species (% A.F.S) designates the percentage of a morphotype compared to the totality of the spores.

$$A. F. S = \left( \frac{N_s}{N_t} \right) * 100$$

N<sub>s</sub> : Number of isolated spores of the species x.

N<sub>t</sub> : Total number of spores.

### Dry weight of tubers

The dry matter of the tubers was determined by steaming by taking the fresh weight of the departure and proceeding to the drying in the study at 80°C until constant weight.

### Nitrogen content of tubers

The determination of nitrogen was carried out by the Kjeldhal method, which consists of the mineralization of organic matter under heat with concentrated sulfuric acid (H<sub>2</sub>SO<sub>4</sub>) in the presence of catalysts (CuSO<sub>4</sub> and K<sub>2</sub>SO<sub>4</sub>).

The diammonium sulfate is then displaced from its soda salt, distilled by steam distillation and collected in a known amount of excess hydrochloric acid. The unreacted hydrochloric acid is measured back with sodium hydroxide.

### Analysis of mineral elements

The analysis of mineral elements was carried out by X-ray fluorescence spectrometry (EDX 7000 Shimadzu); the chemical elements present in the samples are represented by their mass concentration.

### Statistical analysis

Data presented as means of 3 replicates ± standard deviation were analyzed by one-factor ANOVA test using SPSS 21.0 software for Windows.

### Conclusion

We conclude that arbuscular mycorrhizal fungi can be used as a promising approach for high quality and quantity of potato production. Using commercial inoculum and natural inoculum isolated from Guercif-Morocco soil; we observed an increase in yield and biomass, in addition to the improvement of tuber mineral content. Furthermore, the results obtained with the natural inoculum (T3) exceeded those recorded for the commercial inoculum. These results shows the potential of this natural inoculum and the need for the development and improvement of such preparation and to test it on other crops.

### Acknowledgments

I thank Said Lachgar for technical assistance and Hicham Labazi for editing the manuscript.

### References

- AL-Shmary RF and AL-Taey DKA (2020) The response of two potato cultivars to organic and bio-fertilization and their interaction with chemical fertilization on some growth and yield traits International Journal of Psychosocial Rehabilitation. 24(6) : 6774-6780.
- Amir H, Renard A (2003) Etude microbiologique générale de quelques sols de forêts sclérophylles de Nouvelle-Calédonie: Statuts des mycorhizes à arbuscules. PCFS-UNC CP. 54, 22.

- Béreau M, Garbaye J (1994) First observations on the root morphology and symbioses of 21 major tree species in the primary tropical rain forest of French Guyana. *Ann For Sci.* 51: 407–416. <https://doi.org/10.1051/forest:19940406>.
- Bhandari P, Garg N (2017) Dynamics of arbuscular mycorrhizal symbiosis and its role in nutrient acquisition: an overview. *Mycorrhiza- Nutrient Uptake, Biocontrol.* Springer International Publishing, Ecorestoration Cham. pp 21–43.
- Bharadwaj D P, Lundquist P, and Alströma S (2007) Impact of plant species grown as monocultures on sporulation and root colonization by native arbuscular mycorrhizal fungi in potato. *Appl Soil Ecol.* 35:213-225. DOI:10.1016/j.apsoil.2006.04.003.
- Boussageon R, van Tuinen D, Lapadatescu C et al (2023) Effects of field inoculation of potato tubers with the arbuscular mycorrhizal fungus *Rhizophagus irregularis* DAOM 197198 are cultivar dependent. *Symbiosis.* 89: 213–226. <https://doi.org/10.1007/s13199-023-00908-w>.
- Chafai W, EL Gabardi S, Douira A and Khalid A (2022) Diversity and mycorrhizal potential of arbuscular mycorrhizal fungi in two natural soils in the eastern region of Morocco. *Asian J Agric Biol.* 202102101. DOI: 10.35495/ajab.2021.02.101.
- Chliyah M, Ouazzani Touhami A, Filali-Maltouf A, El Modafar C, MoukhliA, Oukabli A, Benkirane R, Douira A (2014) Effect of a composite endomycorrhizal inoculum on the growth of olive trees under nurseries conditions in Morocco. *Int J Pure App Biosci.* 2: 1-14.
- Cobb AB, Duell EB, Smith FA, Smith SE (2021) Utilising mycorrhizal responses to guide selective breeding for agricultural sustainability. *Plants People Planet.* 3: 578-587.
- Derkowska E, Sas Paszt L, Sumorok B, Szwonek E, Gluszek S (2008) The influence of mycorrhization and organic mulches on mycorrhizal frequency in apple and strawberry roots. *J Fruit Ornam Plant Res.* 16:227- 242.
- Ettlili S, Labidi S, Khiari B, Jerbi M, Ben Alaya A, Djébalı N (2022) Reduced nitrogen and phosphorus fertilization combined with mycorrhizal inoculation enhance potato yield and soil mineral fertility. *J Oasis Agric Sustain.* 4(4), 20–29. <https://doi.org/10.56027/JOASD.212022>.
- FAO (2015) FAOSTAT. Food and Agriculture Organization of the United Nations.
- Ferrer RL, Herrera RA (1980) El género *Gigaspora* Gerdemann et Trappe (Endogonaceae) en Cuba. *Revista Jard Bot Nac Habana.* 1(1): 43–66.
- Gerdemann J W and Nicolson T H (1963) Spores of mycorrhizal Endogone species extracted from soil by wet sieving and decanting. *Trans Brit Mycol Soc.* 46(2), 235-246. [http://dx.doi.org/10.1016/S0007-1536\(63\)80079-0](http://dx.doi.org/10.1016/S0007-1536(63)80079-0).
- Hall IR (1984) Taxonomy of VA mycorrhizal fungi. In: Powell CL, Bhagyaraj DJ, editors. *VA Mycorrhiza.* Boca Raton. FL: CRC Press, 57–94.
- Hart M, Ehret DL, Krumbein A, Leung C, Murch S, Turi C and Franken P (2015) Inoculation with arbuscular mycorrhizal fungi improves the nutritional value of tomatoes. *Mycorrhiza.* 25 (5), 359-376.
- Hassan AA (2003) *Potato.* Dar-AL-Arabiya Publication. Cairo. Egypt. 198.
- Hepper CM, Azco’n-Aguilar C, Rosendahl S and Sen R (1988) Competition between three species of *Glomus* used as spatially separated introduced and indigenous mycorrhizal inocula for leek (*Allium porrum* L.). *New Phytol.* 110:207–215. <https://doi.org/10.1111/j.1469-8137.1988.tb00254.x>.
- Higo M, Azuma M, Kamiyoshihara Y, Kanda A, Tatewaki Y, Isobe K (2020) Impact of Phosphorus Fertilization on Tomato Growth and Arbuscular Mycorrhizal Fungal Communities. *Microorganisms.* 8 (2):178. <https://doi.org/10.3390/microorganisms8020178>
- INVAM (2017) International Culture Collection of (Vesicular) Arbuscular Micorrhizal Fungi, Arbuscular Mycorrhizal Fungi, Diperoleh dari West Virginia University.
- Jansa J, Mozafar A, Kuhn G, Anken T, Ruh R, Sanders IR, Frossard E (2003) Soil tillage affects the community structure of mycorrhizal fungi in maize roots. *Ecol Appl.* 13: 1164-1176. [https://doi.org/10.1890/1051-0761\(2003\)13\[1164:STATCS\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2003)13[1164:STATCS]2.0.CO;2).
- Jansa J, Smith FA and Smith SE (2008) Are there benefits of simultaneous root colonization by different arbuscular mycorrhizal fungi. *New Phytol.* 177:779–789. DOI: 10.1111/j.1469-8137.2007.02294.x.
- Khoshnevisan B, Rafieei S, Omid M, Mousazadeh H (2013) Comparison of GHG emission of efficient and inefficient potato producers based on data envelopment analysis. *J Agric Eng Biotec.* 1(3): 81-88. DOI:10.18005/JAEB0103005.
- Koide RT (2000) Functional complementarity in the arbuscular mycorrhizal symbiosis. *New Phytol.* 147: 233– 235. <https://doi.org/10.1046/j.1469-8137.2000.00710.x>.
- Kumar R and Keshar dev (2017) Effects of Chemical Fertilizers on Human Health and Environment: A Review. *International Advanced Research Journal in Science. Eng Technol.* 4: 2393-8021.
- Lombardo S, Abbate C, Pandino G, Parisi B, Scavo A and Mauromicale G (2020) Productive and physiological response of organic potato grown under highly calcareous soils to fertilization and mycorrhization management. *J Agron.* 10(8): 1200. <https://doi.org/10.3390/agronomy10081200>.
- Morton JB, Benny GL (1990) Revised classification of arbuscular mycorrhizal fungi (Zygomycetes): a new order, Glomales, two new suborders, Glomineae and Gigasporineae, and two new families, Acaulosporaceae and Gigasporaceae, with an emendation of Glomaceae. *Mycotaxon.* 37: 471–491.
- Ndiaye F, Diop TA, Sy MO, Manga AGB, Sow HA, Ba T (2005) Controlled mycorrhization of micropropagated potato plants (*Solanum tuberosum* L.). *J Agric Food Res.* 1:33-37.
- Ortas I, Ustuner O (2014) Determination of different growth media and various mycorrhizae species on citrus growth and nutrient uptake. *Sci Hortic.* 166: 84-90. <https://doi.org/10.1016/j.scienta.2013.12.014>.
- Plenchette C, Clermont-Dauphin C, Meynard JM, Fortin JA (2005) Managing arbuscular mycorrhizal fungi in cropping systems. *Can J Plant Sci.* 85: 31-40. <https://doi.org/10.4141/P03-159>.
- Pearson JN, Abbott LK, Jasper DA (1993) Mediation of competition between two colonizing VA mycorrhizal fungi by host plants. *New Phytol.* 123:93–98. <https://doi.org/10.1111/j.1469-8137.1993.tb04534.x>.
- Püschel D, Bitterlich M, Rydlová J, Jansa J (2021) Drought accentuates the role of mycorrhiza in phosphorus uptake. *Soil Biol Biochem.* 157: 108243.
- Ravnskov S, Larsen J, Jakobsen I (2002) Phosphorus uptake of an arbuscular mycorrhizal fungus is not effected by the biocontrol bacterium *Burkholderia cepacia*. *Soil Biol Biochem.* 34: 1875–1881. doi: 10.1016/S0038-0717(02)00201-8
- Ruytinx J, Kafe A, Usman M, Coninx L, Zimmermann SD, Garcia K (2020) Micronutrient transport in mycorrhizal symbiosis; zinc steals the show. *Fungal Biol Rev.* 34:1–9. DOI:10.1016/j.fbr.2019.09.001.
- Sánchez-Bel P, Sanmartín N, Pastor V, Mateu D, Cerezo M, Vidal-Albalat A, Pastor-Fernández J, Pozo MJ, Flors V (2018)



- Mycorrhizal Tomato Plants Fine Tunes the Growth-defence Balance upon N Depleted Root Environments. *Plant Cell Environ.* 41 (2): 406–420. DOI: 10.1111/pce.13105
- Sánchez-Romera B, Ruiz-Lozano JM, Zamarreño ÁM, et al (2016) Arbuscular mycorrhizal symbiosis and methyl jasmonate avoid the inhibition of root hydraulic conductivity caused by drought. *Mycorrhiza.* 26:111–122. DOI: 10.1007/s00572-015-0650-7.
- Schenck NC, Pérez Y (1987) *Manual for the identification of VA mycorrhizal fungi*, first. ed. Gainesville, FL: University of Florida, School of Forest Resources and Conservation. 286 p.
- Schenck NC, Smith G (1982) Additional new and unreported species of mycorrhizal fungi (Endogonaceae) from Florida. *Mycologia*, 74(1): 77–92. <https://doi.org/10.1080/00275514.1982.12021472>.
- Selmaoui K, Artib M, Semane F, El Gabardi S, Hibilik N, El aymani I, Chliyah M, Mouria A, Ouazzani Touhami A, Benkirane R, Douira A (2017) Diversity of endomycorrhizal fungi (AMF) in the rhizosphere of sugarcane (*Saccharum officinarum*) grown in Morocco. *Int J Recent Sci Res*, 8(2): 15753-15761.
- Sghir F, Touati J, Chliyah M, Ouazzani Touhami A, Filali-Maltouf A, El Modafar C, Moukhli A, Oukabli A, Benkirane R, Douira A (2014) Diversity of arbuscular mycorrhizal fungi in the rhizosphere of date palm tree (*Phoenix dactylifera*) in Tafilalet and Zagora regions (Morocco), *Int J Pure App Biosci*, 2 (6): 1-11.
- Tahat MM, Kamaruzaman S, Radziah O, Kadir J, Masdek HN (2008) Response of (*Lycopersicon esculentum* Mill.) to different arbuscular mycorrhizal fungi species. *Asian J Plant Sci*, 7: 479–484. DOI: 10.3923/ajps.2008.479.484.
- Talbi Z, El Asri A, Touati J, Chliyah M, Ait aguil F, Selmaoui K, Sghir F, Ouazzani Touhami A, Benkirane R, Douira A (2015) Morphological characterization and diversity of endomycorrhizae in the rhizosphere of Carob tree (*Ceratonia siliqua*) in Morocco. *IJBLS*, 3(1):196-211.
- Thirkell TJ, Grimmer M, James L, Pastok D, Allary T, Elliott A, Paveley N, Daniell T, Field KJ (2022) Variation in mycorrhizal growth response among a spring Wheat mapping population shows potential to breed for symbiotic benefit. *Food Energy Secur.* 11, e370.
- Tohidi-Moghaddam H, Sani B, Ghooshchi F (2004) The effect of nitrogen fixing and phosphate solubilizing microorganism on some quantitative parameters on soybean from sustainable agricultural point of views. *Proceeding of 8th Agronomy and Plant Breeding Congress of Iran*, Guilan University, Iran.
- Trouvelot A, Kough JL (1986) Measurement of the VA mycorrhization rate of a root system. Search for estimation methods with functional significance. In *Physiological and genetical aspects of mycorrhizae*. Gianinazzi-Pearson V, Gianinazzi S (Eds.), INRA édition, Paris, 217-221.
- Vierheilig H, Garcia-Garrido JM, Wyss U, Piche Y (2000) Systemic suppression of mycorrhizal colonization of barley roots already colonized by AM fungi. *Soil Biol Biochem*, 32:589–595. [https://doi.org/10.1016/S0038-0717\(99\)00155-8](https://doi.org/10.1016/S0038-0717(99)00155-8).
- Vierheilig H (2004) Further root colonization by arbuscular mycorrhizal fungi in already mycorrhizal plants is suppressed after a critical level of root colonization. *J Plant Physiol.* 161:339–341. <https://doi.org/10.1078/0176-1617-01097>.
- Vosatka M, Gryndker M (2000) Response of micropropagation potatoes transplanted to peat media to post vitro inoculation with arbuscular mycorrhizal fungi and soil bacteria. *Appl Soil Ecol.* 15: 145-152. DOI: 10.5897/AJMRx12.005.
- Wu F, Wang W, Ma Y, Liu Y, Ma X, An L, Feng H (2013) Prospect of beneficial microorganisms applied in potato cultivation for sustainable agriculture. *Afr J Microbiol Res.* 7: 2150– 2158. DOI: 10.5897/AJMRx12.005.
- Walker C, Mize CW, Mc Nabb HS (1982) Populations of endogonaceous fungi at two localities in central Iowa. *Canad J Bot.* 60(12): 2518–2529. <https://doi.org/10.1139/b82-305>.
- Zhong Qun H, Chao Xing H, Zhibin Z, Zhirong Z, Huai Song W (2007) Changes in antioxidative enzymes and cell membrane osmosis in tomato colonized by arbuscular mycorrhizae under NaCl stress. *Colloids Surf B.* 59:128–133. doi: 10.1016/j.colsurfb.2007.04.023. Epub 2007 May 5.