

Foliar application of chitosan and plant probiotic bacteria influencing the growth, productivity and bulb storage life of onion

Tithy Ray¹, Joydeb Gomasta^{*1}, Jahidul Hassan¹, Md. Shamim Hossain² and Emrul Kayesh^{*1}

¹Department of Horticulture, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh

²Department of Entomology, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh

*Corresponding author: ekayeshrt@bsmrau.edu.bd; joydeb@bsmrau.edu.bd

Abstract

Onion having several health promoting phytochemical contents is an important spices in most of the countries of the world as well as vegetables in particular and chitosan and plant probiotic bacteria are few proven natural growth enhancing agents. Here, three bio-stimulants namely Chitosan (100 and 200 ppm), *Paraburkholderia* BRRh-4 bacteria (1.5×10^9 CFU mL⁻¹) and Clybio (0.1%) in singly as well as combining one another were applied at twelve different treatment combinations including control with the aim of investigating the below and above ground vegetative growth, bulb productivity and post-harvest storage performance of onion cv. "Taherpuri". The experiment was set up in Randomized Complete Block Design (RCBD) maintaining three replicates and conducted during November 2019 to June 2020 where the bio-stimulant formulations were sprayed in the foliage at 20, 40 and 60 days after transplanting (DAT). Reproductive behaviors of the onion cultivar differed significantly among the treatments and those were the resultants of statistically varied vegetative performances recorded at 30, 60 and 90 DAT. Chitosan @ 100 ppm + Clybio (0.1%) + *Paraburkholderia* BRRh-4 (1.5×10^9 CFU mL⁻¹) combination exhibited statistical superiority over the other treatments in terms of plant height (53.230 cm), number of leaves (10.13/plant), leaf length (41.20 cm) and diameter (10.33 mm), number of roots (32.53/plant), root length (12.23 cm), neck and bulb diameter (11.20 mm and 4.83 cm, respectively) and bulbing ratio (0.23) at 90 DAT. While, shoot and root growth didn't vary significantly at earlier stages (at 30 DAT). Again, the highest yield (3.36 kg/plot and 10.45 t/ha) was also obtained from Chitosan @ 100 ppm + Clybio (0.1%) + *Paraburkholderia* BRRh-4 (1.5×10^9 CFU mL⁻¹) treatment. The combination of chitosan and probiotic bacteria had significant influence on post-harvest storage qualities of onion where the longest shelf life/marketable life (8.17 months) with the lowest physiological weight loss and bulb rotting at 90 days after harvest was administered in Chitosan @ 100 ppm + Clybio (0.1%) + *Paraburkholderia* BRRh-4 (1.5×10^9 CFU mL⁻¹) treatment. Besides the best treatment, Chitosan @ 100 ppm + Clybio (0.1%) and Chitosan @ 100 ppm + *Paraburkholderia* BRRh-4 (1.5×10^9 CFU mL⁻¹) produced statistically superior results over control for onion growth and development. Therefore, chitosan and plant probiotic bacteria can be judiciously used for quality onion production in the warm and humid areas of the world.

Keywords: Onion (*Allium cepa* L.); shrimp shell chitosan; *Paraburkholderia* BRRh-4; clybio; root and shoot growth; bulb yield; storage quality.

Abbreviations: CFU_Colony forming unit, DAH_Days after harvest, DAT_Days after transplanting, PLW_Physiological loss in weight, RH_Relative humidity, T-Max_Maximum temperature, T-Min_Minimum temperature

Introduction

Onion (*Allium cepa*) is the most important member of Alliace family which is used worldwide as culinary and therapeutic spice and vegetable as well. It is an crucial ingredient in many Asian and African sauces and is primarily grown locally, with Egypt being the continent's pioneer producer (Kuete, 2017). The crop has a great economic importance and is the second most important spice crop in the world. Onion bulb is a good source of carbohydrates, proteins, vitamins (B₆, C), minerals (phosphorus, calcium, magnesium, iron, manganese, sulfur), amino acids and antioxidants. In addition, a variety of secondary metabolites have been identified in onion species such as flavonoids, phenolics, phytosterols, and saponins (Ware, 2017; Marrelli et al., 2019) that are known to diminish cardiovascular ailment, diabetes and antibacterial, antiviral, anti-allergenic.

The green leaves and lowering stalks are also edible and popular. Onion is famous for its characteristics flavor and widely used to increase the taste of foods like gravies, soups, stew stuffing, fried fish and meat but can also be eaten raw or used to make pickles and chutneys (Rashid et al., 2015). In the south-Asian and African countries onion is a high-value spice crop. Households' daily kitchen chores and restaurants as well as the industries demand quality onion throughout the year. But most of the high onion consuming countries fall short of quality onion production in adequate amount. In order to increase crop productivity and improve soil fertility, the farmers indiscriminately use various inorganic fertilizers and crop boosters, which might cause several environmental and health damages (Parr et al., 1991). Also, the intensive use of inorganic fertilizers without

the addition of organic matter leads to poor soil strata, which is more sensitive to wind and rain erosion. To overcome the above problems, environmental advocates and many investigators recommended the use of beneficial soil micro-organisms and products derived from biological transformations. Chitosan, clybio and plant probiotic bacteria might be some key inclusive of such eco-friendly growth booster of crops.

Chitosan, a polycationic polymer of β -1,4, linked D-glucosamine chemically derived from crustaceans and soluble in organic acids is one of a range of natural compounds that have been shown to be effective in decreasing disease incidence in crops (Jitareerat et al., 2007). It is considered environmentally safe for various agricultural uses as it is easily biodegradable, biocompatible, and non-toxic to human beings (Basit et al., 2020). Chitosan and its derivatives have been reported to act as a plant growth regulator and are considered to elicit natural defense responses in plants and have been used as a natural compound to control pre-and post-harvest pathogenic diseases as well as abiotic stresses like drought, salinity, etc. (Uthairatanakij et al., 2007). However, many investigators reported that using chitosan as a foliar spray increased the vegetative growth, yield, and biochemical contents of vegetable crops (El-Miniawy et al., 2013). On the other hand, plant probiotic bacteria are naturally occurring plant-associated microorganisms that enhance the growth of the host plants including yield and may suppress diseases when applied in adequate amounts. They provide beneficial effects to host plants through the production of phytohormones, antibiotics, and lytic enzymes, fixation of atmospheric nitrogen, solubilization of soil mineral nutrients, and induction of systemic resistance in the host plants (Islam and Hossain, 2012). Clybio formulation is the mixture of three probiotics namely yeast fungus, *Bacillus natto* and *Lactobacillus*. Yeast is a natural, safe source of biofertilizer which is typically applied on soil or foliar application on crops improved growth characters and yields of vegetable crops while *Bacillus* and *Lactobacillus* can act as the soil probiotic making it a perfect medium for organic farming to enhance plant growth and protect against disease causing organisms (Ikeda et al., 2013). However, scientific information is meager on the influence of combined applications of chitosan, plant probiotic bacteria and clybio formulation on yield, yield contributing attributes and post-harvest storage life of onion. Beside yield enhancement, these growth regulators can add valuable anti-oxidants and mineral elements to the produces. Considering the above facts, the present investigation was undertaken to observe the influence of chitosan and plant probiotic bacteria on the vegetative and reproductive performance including bulb yield and physical post-harvest attributes of onion in Bangladesh condition.

Results

Effect on vegetative growth characters

Plant height

Significant variations were noticed among the treatments at 30, 60 and 90 days after transplanting (DAT) for plant height in onion and an increase in plant height was noted with the foliar application of chitosan alone or in combination with *Paraburkholderia* and clybio than control plants (Figure 1). At 30 and 60 DAT, the tallest plant (33.07 cm and 47.33 cm, respectively) was obtained from Chitosan @ 200 ppm +

Paraburkholderia combination (T_{11}) and Chitosan @ 200 ppm + Clybio combination (T_{10}) treatments, respectively having been statistically similar to that of all other treatments except T_1 treatments which exhibited the shortest plant throughout the season (26.68 cm, 42.60 cm and 44.13 cm at 30, 60 and 90 DAT, respectively). While at 90 DAT the longest plant (53.20 cm) was recorded in Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8) which was statistically succeeded by T_9 , T_{10} , and T_{11} treatments.

Number of leaves per plant

Number of leaves per plant varied significantly in response to various treatments at 30 and 90 DAT, whereas at 60 DAT number of leaves per plant didn't differ statistically (Table 1). At 30 DAT, maximum number of leaves per plant (3.47) was counted in Chitosan @ 200 ppm + *Paraburkholderia* combination (T_{11}) which was closely followed by T_2 , T_3 and T_8 treatments. Although there was no statistical difference in the number of leaves per plant among the treatments, but numerically the highest number of leaves per plant (5.93) was recorded in Chitosan @ 200 ppm + Clybio (T_{10}) treatment at 60 DAT. At 90 DAT, statistically maximum number of leaves per plant (10.13) was registered in Chitosan @ 100 ppm + Clybio + *Paraburkholderia* (T_8) treatment followed by T_7 and T_{10} treatments. Control plants produced minimum number of leaves per plant (3.07, 5.47 and 7.73, respectively) at all the DATs.

Leaf growth

Leaf length and leaf width had significant variation in response to different chitosan and probiotic treatments at different DATs (Table 1). The highest leaf length and leaf diameter was estimated 31.70 cm and 4.93 mm, respectively in plants treated with Chitosan @ 200 ppm + *Paraburkholderia* combination (T_{11}) at 30 DAT having statistical similarity to most other treatments except T_1 , T_7 and T_9 for leaf length and T_1 and T_{12} for leaf width. Again, at 60 and 90 DAT maximum leaf length and leaf width (41.20 cm and 46.87 cm and 6.80 mm and 10.33 mm, respectively) was resulted in the plants sprayed with Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8). Statistically inferior growth in terms of leaf length and leaf width was noticed in untreated plants.

Root growth

Onion root number and root length were also significantly influenced by different levels of treatments (Table 2). Although there was no statistically significant difference in plant root number among the treatments, but numerically higher and lower number of roots (21.73 and 18.13) was counted in plants treated with Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8) and control treatment (T_1) at 30 DAT. Again, at 60 and 90 DAT, maximum number of roots (24.53/plant and 32.53/plant) was recorded from Chitosan @ 0 ppm + *Paraburkholderia* (T_3) and Chitosan @ 100 ppm + Clybio + *Paraburkholderia* (T_8) treatments, respectively. On the other hand, at 60 and 90 DAT, the lengthiest root (8.71 cm and 12.23 cm, respectively) was estimated in plants treated with Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8) which was statistically followed by T_{11} and T_{12} treatments. Whereas, root number as well as root length at 60 and 90 DAT was noted minimum (19.07 and 25.53 and 5.87 cm and 8.75 cm, respectively) in control plants.

Neck diameter

Neck diameter differed significantly in response to different treatments at 30, 60 and 90 days after transplanting of onion seedlings (Table 3). Therefore, Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8) performed the best in terms of neck diameter at all the DATs (3.97 mm, 8.33 mm and 11.20 mm, respectively) and statistically and numerically identical neck diameter was also revealed by Chitosan @ 100 ppm + Clybio treatment (T_6) at both 60 and 90 DAT. Furthermore, neck diameter was observed to be statistically unique at 60 and 90 DAT for all the treatments except control which exhibited minimum diameter (2.20 mm, 6.80 mm and 9.73 mm, respectively) at 30, 60 and 90 DAT.

Bulb diameter

Bulb diameter was positively influenced and varied significantly due to different treatments (Table 3). Statistically maximum and numerically same bulb diameter (1.33 cm) was registered in Chitosan @ 200 ppm + Clybio (T_{10}) and Chitosan @ 200 ppm + Clybio + *Paraburkholderia* (T_{12}) combinations at 30 DAT which was statistically similar to all other treatments except T_2 and T_1 treatments. While, at 60 DAT, the highest bulb diameter (2.56 cm) was recorded from the treatment Chitosan @ 200 ppm + Clybio (T_{10}) which was statistically identical to that of all others except T_1 treatment. Again, bulb diameter at 90 DAT was noticed the highest (4.83 cm) in Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8) which was similar to T_5 , T_6 , T_7 and T_{10} treatments. But minimum bulb diameter (1.13 cm, 2.03 cm and 3.28 cm, respectively) was observed in control (T_1) treatment at all the DATs.

Bulbing ratio

At 30 DAT, significantly maximum bulbing ratio (0.30) was observed in Chitosan @ 100 ppm + Clybio + *Paraburkholderia* treatment (T_8) followed by T_6 , T_{10} , T_{11} and T_{12} treatments, while minimum bulbing ratio (0.20) was noted in control (T_1) treatment (Table 3). On the other hand, the highest bulbing ratio (0.38) at 60 DAT was obtained by foliar spraying the onion plants with Chitosan @ 200 ppm (T_9) treatment and the lowest bulbing ratio (0.28) was recorded in control (T_1) and Chitosan @ 0 ppm + *Paraburkholderia* (T_3). Furthermore, the lowest bulbing ratio (0.23) at 90 DAT was obtained from Chitosan @ 100 ppm + Clybio + *Paraburkholderia* combination (T_8) which was statistically similar to all other treatments except T_1 and T_{12} treatment. Thus, foliar application of different concentrations of chitosan and probiotic bacteria treatment on onion plants produced the lowest bulbing ratio at 90 DAT.

Effect on yield and yield contributing characters

Splitting of bulb

Splitting of bulbs showed a significant response to different treatments (Figure 2). Maximum percentage of splitting bulb was calculated in T_{11} (11.08%) treatment followed by T_3 (10.82%), T_7 (11.05%), T_{10} (10.54%) which was statistically similar to the other treatments and the minimum was recorded in control T_1 (5.69%) which was statistically different from other treatments.

Single bulb weight and bulb yield

Single bulb weight showed a significant response to different treatments at 30, 60, and 90 DAT (Table 4). At 30 DAT, the heaviest bulb was produced by T_8 (1.96 g) treatment and the lightest bulb was produced by T_1 (1.32 g) treatment. On the other hand, the highest bulb weight was observed from T_8 (11.18 g) treatment at 60 DAT. Treatment T_{12} (10.70 g) produced the second-highest bulb weight and the lowest bulb weight was found in T_1 (7.68 g) treatment at 60 DAT which was statistically different from all other treatments. Furthermore, the maximum individual bulb weight was recorded from T_8 (42.25 g) treatment followed by T_6 (40.64 g), T_{12} (39.57 g) and the minimum individual bulb weight was found from T_1 (30.70 g) at 90 DAT. Following this trend, significantly maximum yield was recorded from the treatment T_8 (3.36 kg/plot and 10.45 t/ha) which was statistically similar to T_6 (3.08 kg/plot and 9.50 t/ha) treatment, while the minimum yield per plot was found in T_1 (2.24 kg/plot and 6.89 t/ha) treatment.

Effect on post-harvest quality attributes

Physiological weight loss

Significant ($p \leq 0.05$) variation was observed in the percentage of weight loss according to the different treatments in various storage durations (10-90 days) (Table 5). Weight loss was noticed maximum in control (T_1) treatment throughout the storage period of onion. On the other hand, the minimum physiological weight loss was recorded from T_8 followed by T_5 , T_6 , T_9 , T_{12} treatment at 90 days of storage.

Rotten bulbs (%)

The percentage of rotten bulbs of onion at different periods of storage was significantly influenced by different treatments (Table 6). The maximum percentage of rotten bulbs (15.14 %) after 90 days of storage was noted in T_1 which was statistically different from the other treatments, whereas the lowest percentage of rotten bulbs was observed in T_8 (5.66 %) treatment followed by T_5 , T_6 , T_9 , T_{12} treatments.

Shelf life

Shelf life of onions was greatly influenced and statistically significant ($p \leq 0.05$) by different treatments (Figure 3). The longest shelf life was found in T_8 (8.17 months) treatment, while the lowest shelf life was observed in T_1 (3.50 months) control. However, shelf life of onions was significantly enhanced in T_5 (6.83 months), T_6 (6.90 months), T_7 (6.14 months), T_{10} (6.50 months), and T_{12} (6.80 months) treatment, respectively.

Discussions

Onion is one of the most important spices in terms of its economic value and also valuable for containing several Many biologically active substances, including phenolic acids, thiosulfinates and flavonoids with proposed health-related properties including antidiabetic, antimicrobial, cardiovascular, anticancer, antioxidant effects, etc. and this important spice crop requires balanced supply of nutrient inputs, chemical as well as naturally occurring growth promoters, for successful growth, development and yield. In the present research, foliar application of chitosan @ 100 ppm when combined with both of clybio @ 0.1% and *Paraburkholderia* BRRh-4 bacteria @ 1.5×10^9 CFU mL⁻¹ or separately with each of the two at early vegetative growth phases significantly improved root and shoot growth, neck

and bulb diameter resulting in the statistically lowest bulbing ratio at 90 days after transplanting (Figure 4) compared to the control. Again, the superior vegetative growth with chitosan and probiotic bacteria ultimately improved the bulb yield and post-harvest qualities of onion with extended shelf life (Figure 5).

Shrimp shell chitosan and plant probiotic bacteria have been now-a-days used in crop production as non-chemical growth and yield promoting agents as these increase the availability and uptake of water and essential nutrients through adjusting cell osmotic pressure and reducing the accumulation of harmful free radicals by increasing the enzymatic and other functional activities (Abdel-Mawgoud et al., 2010; Garcia-Seco et al., 2015; Yildirim et al., 2015; Emami Bistgani et al., 2017; Jiménez-Gómez et al., 2017; Hassnain et al., 2020; Chakraborty and Islam, 2022). Plant growth-promoting bacteria improve plant growth parameters by increasing the levels of auxins and cytokinins and decreasing the levels of ethylene and abscisic acid in the plants, which result in enhanced cell division and elongation of plants (Ruzzi and Aroca, 2015). Mondal et al. (2012) opined that the application of chitosan increased the key enzyme activities of nitrogen metabolism (nitrate reductase, glutamine synthetase, and protease) and improved the transportation of nitrogen (N) in the functional leaves as well as increased photosynthesis which enhanced plant growth and development. In the present investigation, chitosan, *Paraburkholderia* and clybio spray improves the vegetative growth indicating increased number of leaves, leaf length and diameter, number of roots and root length in onion over control. Alike this experiment, it is reported that foliar application of chitosan and rhizobacteria alone significantly increased plant growth and development in onion (Dilpreet et al., 2016; Kumar et al., 2017; Sutariati et al., 2019), garlic (Ahmed and Farm, 2015), potato (Shaheen et al., 2019) and pepper (Shrestha et al., 2014). Mukta et al. (2017) and Rahman et al. (2018a,b) also reported that foliar application of chitosan, *Bacillus* and *Paraburkholderia* improved root and shoot growth in treated plants compared to non-treated plants in strawberry. *Bacillus* isolates also have involvement in the modulation of plant development through the production of phytohormones, efficient in producing auxin that might stimulate root proliferation (Bashan and de-Bashan, 2005; Jiménez-Gómez et al., 2016) and nutrient uptake (Tsavkelova et al., 2006) resulting in the vegetative growth of plants as occurred here in the present experiment with onion upon application of clybio consisting of yeast fungus, *Bacillus natto* and *Lactobacillus*. The valuable components of yeast, such as cytokinins, can stimulate cell division and elongation, as well as the synthesis of proteins and nucleic acids, and can increase mineral nutrients and can potentially increase crop growth (Fawzy et al., 2012; Matter and Abou-Sreya, 2016). Lactic acid bacteria strains like *Lactobacillus* are mostly studied as probiotics organisms because they produced various antibacterial compounds, such as indole-3-acetic acid (IAA), lactic acid, hydrogen peroxide, several bacteriocins, and even antifungal compounds (Hamed et al., 2011; Axel et al., 2012; Giassi et al., 2016), which is responsible for plant growth promotion (Mohite, 2013; Shrestha et al., 2014). Reetha et al. (2014) and Tinna et al. (2020) also noted that Rhizobacteria isolates of *Pseudomonas fluorescens* and *Bacillus subtilis* significantly produced higher shoot lengths compared to control plants of onion. Enhanced bulb diameter and consequently low bulbing ratio with the use of

the rhizospheric bacteria *Azotobacter chroococcum*, *Pseudomonas fluorescens*, and *Bacillus subtilis* was reported by Čolo et al. (2014). Present investigation revealed that, foliar application of chitosan and probiotic bacteria produced the lowest bulbing ratio at harvest and that low bulbing ratios are desirable for onion growers (Abdel-Moneim et al., 2015). Splitting of bulb was also noticed in some cases which might be due to fluctuations in day and night temperature during the bulb formation (Steer, 1980) rather than the effects of treatments.

Again, the increase in yield might be due to significantly higher plant height, leaf growth and root growth with combined application of chitosan and probiotic bacteria inoculation leading to the higher synthesis of photosynthates and their better translocation to the sink, as the rate of photosynthesis is significantly correlated with the growth of onion. Čolo et al. (2014) observed that inoculation of onion with *Azotobacter chroococcum*, *Bacillus subtilis* and the mixture of the inoculants significantly increased bulb weight and yield in onion. Similarly Afify et al. (2018) examined that the microbial inoculation combination with chemical fertilizers improves the growth and productivity of plants and increases bulb weights and total bulb yield of onion. Identical observations on enhanced yield and yield contributing characters was noted in several crops through foliar application of chitosan [Sultana et al. (2017) and Parvin et al. (2019) in eggplant and tomato, Rahman et al. (2018a) and Soppelsa et al. (2019) in strawberry, Rodríguez et al. (2017) in potato, Mahmood et al. (2017) in bell pepper, Mehehub et al. (2019) in litchi], plant probiotic bacteria [Karlidag et al. (2010) and Rahman et al. (2018b) in strawberry, Yildirim et al. (2015) in cabbage, Garcia-Seco et al. (2015) in blackberry] and clybio [Akter et al. (2021) in spinach, Uddin et al. (2021) in strawberry].

Along with higher yield, chitosan, *Paraburkholderia* and clybio combination enhanced the storability of onion through by minimizing physiological weight loss and rotten bulb percentages which might be resultant of high dry matter accumulation in onion bulb as a function of extended photosynthetic CO₂ fixation in leaves and its translocation to the bulb in chitosan and probiotic bacteria inoculated plants. Besides, chitosan and plant probiotic bacteria are organic growth stimulants that extend the storage through the reduction of respiration rate and water loss and prevent the colonization of pathogen, thereby improves the immune system of the onion bulb (Shehata et al., 2012; Gerjes et al., 2020). Kloepper et al. (2004) reported that *Bacillus subtilis* is a biocontrol agent because they protect plant roots from phytopathogenic fungi such as *Fusarium* and *Pythium*. Liu et al. (2016) also opined that chitosan and oligochitosan enhance ginger resistance to rhizome rot in storage. Furthermore, chitosan, as an antimicrobial agent, induces immunity of plants against many plant pathogenic bacteria and fungi and is likely to be linked with prolonging the shelf life of fruits and vegetables in storage conditions (Romanazzi et al., 2017). In the present experiment, similar phenomena might occur and thereby lower percentages of rotting bulbs were obtained from the combination of chitosan with clybio and *Paraburkholderia* BRRh-4 bacteria treatments. Finally, as a result of such inherently developed immune system and post-harvest qualities in the onion bulb significantly reduced the post-harvest loss and enhanced the shelf life up to 8 months after harvest in ambient storage conditions.

Table 1. Effect of chitosan and plant probiotic bacteria on number leaves per plant at different growth stages of onion

Treatment	Number of Leaves /plant			Leaf length (cm)			Leaf diameter (mm)		
	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT
T ₁	3.07c	5.47	7.73e	25.63b	36.40d	40.07d	4.27c	5.30c	7.87e
T ₂	3.40ab	5.67	8.13de	27.90ab	37.76cd	41.53cd	4.60a-c	6.20a-c	8.67d
T ₃	3.40ab	5.60	8.20de	28.60ab	39.27a-d	42.60b-d	4.93a	6.40a-c	8.73cd
T ₄	3.33a-c	5.60	8.67cd	29.93ab	38.27a-d	42.20cd	4.80ab	5.87a-c	8.80cd
T ₅	3.20a-c	5.53	8.53d	31.23a	38.97a-d	41.60cd	4.80ab	5.53bc	9.07b-d
T ₆	3.13bc	5.73	9.33b	27.40ab	38.00b-d	42.00cd	4.60a-c	6.47ab	9.60b
T ₇	3.33a-c	5.80	9.60ab	26.40b	39.47a-d	44.00a-c	4.47a-c	5.73a-c	9.13b-d
T ₈	3.40ab	5.80	10.13a	27.55ab	41.20a	46.87a	4.47a-c	6.80a	10.33a
T ₉	3.2a-c	5.53	9.27bc	26.77b	40.60a-c	45.73a	4.33bc	6.60ab	9.40bc
T ₁₀	3.27a-c	5.93	9.73ab	28.33ab	40.93ab	46.40a	4.53a-c	6.73a	9.20b-d
T ₁₁	3.47a	5.20	9.40b	31.70a	37.83b-d	42.20cd	4.93a	6.27a-c	9.40bc
T ₁₂	3.27a-c	5.53	9.40b	27.57ab	38.77a-d	45.33ab	4.33bc	6.33a-c	9.40bc
LSD _(0.05)	0.297	ns	0.66	4.368	3.136	3.0187	0.5237	1.1374	0.6748
CV (%)	5.34	4.21	4.33	9.13	4.75	4.11	6.74	10.86	4.36

Mean values within a column followed by the same letter do not differ significantly by Fisher's protected LSD test at $p \leq 0.05$, ns= not significant.

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]




Figure 1. Average plant height of onion at 30 DAT (A), 60 DAT (B) and 90 DAT (C) after foliar application of chitosan and plant probiotic bacteria. Vertical bars on the top of the columns represent the standard errors of means of three replicates. Different letters indicate the statistical differences among the treatments at $p \leq 0.05$, as per Fisher's least significant difference test.

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]

Table 2. Effect of chitosan and plant probiotic bacteria on number of roots and root length at different growth stages of onion.

Treatment	Number of roots			Root length (cm)		
	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT
T ₁	18.13	19.07c	25.53e	3.67b	5.87e	8.75e
T ₂	19.13	23.60ab	29.80b-d	3.87ab	6.40d	9.61d
T ₃	19.60	24.53a	30.37b-d	3.97ab	6.50cd	10.05cd
T ₄	18.80	22.20a-c	28.93cd	3.90ab	6.60cd	9.76cd
T ₅	21.13	21.40a-c	30.33b-d	4.40a	6.53cd	10.37c
T ₆	19.00	22.07a-c	30.47a-d	3.30b	6.83b-d	10.04cd
T ₇	19.33	23.47ab	31.13ab	3.67b	6.90bc	10.35c
T ₈	21.73	21.87a-c	32.53a	3.83ab	8.71a	12.23a
T ₉	19.60	21.42a-c	28.67d	3.67b	6.63cd	10.29c
T ₁₀	19.93	23.93ab	30.93a-c	3.33b	6.60cd	10.37c
T ₁₁	19.53	20.93bc	31.33ab	3.53b	7.13b	11.04b
T ₁₂	20.53	22.87ab	31.60ab	3.77ab	7.30b	11.37b
LSD (0.05)	ns	3.4908	2.0909	0.7234	0.4685	0.6197
CV (%)	6.15	9.25	4.1	11.42	4.05	3.54

Mean values within a column followed by the same letter do not differ significantly by Fisher's protected LSD test at $p \leq 0.05$, ns= not significant.

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]




Figure 2. Influence of chitosan and plant probiotic bacteria on splitting of bulb of onion. Vertical bars on the top of the columns represent the standard errors of means of three replicates. Different letters indicate the statistical differences among the treatments at $p \leq 0.05$, as per Fisher's least significant difference test.

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]

Materials and methods

Site details and planting material

The field and laboratory works of experiment was carried out at the research field and laboratory of the Department of Horticulture Bangabandhu Sheikh Mujibur Rahman Agricultural University (BSMRAU), Gazipur-1706, Bangladesh during November 2019 to June 2020. Soil of the experimental field was shallow red-brown terrace soil under the Salna Series (FAO, 1999) in Madhupur Tract (Agroecological Zone 28) having a pH 6.71. The soil had 1.70% organic matter, 0.115% N, 21.35 ppm soluble P and 0.24 meq./100 g soil exchangeable K. The area was characterized by heavy rainfall, excessive humidity, high temperature relatively long days from April to September and scanty rainfall, low humidity, low temperature and a short-day period from October to March. Healthy, insect-pest and pathogen free seeds of the onion cultivar "Taherpuri" were collected from Bangladesh Agricultural Research Institute (BARI), Gazipur, Bangladesh prior to cultivation.

Treatment preparation and application

The treatment of the experiment consists of three types of naturally occurring bio-stimulants namely chitosan polymerase (95% deacetylated) in two separate concentrations (100 ppm and 200 ppm), one plant probiotic bacteria (*Paraburkholderia fungorum* BRRh-4) at 1.5×10^9 CFU mL⁻¹ concentration and clybio (a complex formulation of three probiotics such as yeast fungus, *Bacillus natto*, and *Lactobacillus*) at 0.1% dose in alone and in combinations of any two and all the three types along with no spray. Thus, including control (T₁) a total of twelve different treatment combinations were made namely T₂: Clybio, T₃: *Paraburkholderia*, T₄: Clybio + *Paraburkholderia*, T₅: Chitosan @ 100 ppm, T₆: Chitosan @ 100 ppm + Clybio, T₇: Chitosan @ 100 ppm + *Paraburkholderia*, T₈: Chitosan @ 100 ppm +

Table 3. Effect of chitosan and plant probiotic bacteria on neck and bulb diameter and bulbing ratio at different growth stages of onion.

Treatment	Neck diameter (mm)			Bulb diameter (cm)			Bulbing ratio		
	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT	30 DAT	60 DAT	90 DAT
T ₁	2.20e	6.80c	9.73b	1.13c	2.03b	3.28e	0.20e	0.28b	0.29a
T ₂	2.77d	7.20a-c	10.07ab	1.15bc	2.38a	3.64de	0.24b-d	0.31ab	0.28ab
T ₃	3.00cd	7.93a-c	10.87ab	1.28a	2.52a	3.98cd	0.23c-e	0.28ab	0.28ab
T ₄	2.87cd	7.27a-c	10.87ab	1.27a	2.27ab	3.98cd	0.23de	0.32ab	0.28ab
T ₅	2.80d	6.87bc	10.40ab	1.25ab	2.43a	4.37a-c	0.22de	0.32ab	0.24b
T ₆	3.33bc	8.33a	11.20a	1.23ab	2.30ab	4.60ab	0.27a-c	0.36a	0.24b
T ₇	2.93cd	7.60a-c	10.53ab	1.23ab	2.37ab	4.50a-c	0.24cd	0.36a	0.24b
T ₈	3.97a	8.33a	11.20a	1.31a	2.52a	4.83a	0.30a	0.37a	0.23b
T ₉	3.07cd	8.20ab	11.07ab	1.29a	2.32ab	4.22bc	0.24cd	0.38a	0.27ab
T ₁₀	3.80ab	8.07a-c	10.93ab	1.33a	2.56a	4.43a-c	0.28a	0.32ab	0.25ab
T ₁₁	3.67ab	7.60a-c	10.47ab	1.32a	2.32ab	4.26bc	0.28ab	0.33ab	0.24b
T ₁₂	3.70ab	7.73a-c	10.60ab	1.33a	2.49a	4.22bc	0.28a	0.32ab	0.25ab
LSD _(0.05)	0.5263	1.3956	1.3545	0.1017	0.3227	0.5533	0.0375	0.0705	0.054
CV (%)	9.79	10.76	7.5	4.76	8.02	7.89	8.86	12.67	12.39

Mean values within a column followed by the same letter do not differ significantly by Fisher's protected LSD test at $p \leq 0.05$, ns= not significant.

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]




Figure 3. Effect of chitosan and plant probiotic bacteria on shelf life of onion in storage at room temperature ($25 \pm 1^\circ\text{C}$). Vertical bars on the top of the columns represent the standard errors of means of three replicates. Different letters indicate the statistical differences among the treatments at $p \leq 0.05$, as per Fisher's least significant difference test. [T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]

Table 4. Effect of chitosan and plant probiotic bacteria on single bulb weight at different growth stages and bulb yield (kg/plot and kg/ha) of onion

Treatment	Single bulb weight (g)			Bulb yield	
	30 DAT	60 DAT	90 DAT	kg/plot	kg/ha
T ₁	1.32 c	7.68 d	30.70 f	2.24 d	6.89 d
T ₂	1.41 bc	9.73 bc	32.98 ef	2.90 bc	8.93 bc
T ₃	1.48 bc	9.76 bc	35.21 de	2.61 c	8.04 c
T ₄	1.39 bc	9.43 bc	32.82 ef	2.98 b	9.18 b
T ₅	1.44 bc	9.20 c	34.43 de	3.02 b	9.32 b
T ₆	1.45 bc	9.26 c	40.64 ab	3.08 ab	9.50 ab
T ₇	1.48 bc	9.45 bc	36.51 cd	2.82 bc	8.8 bc
T ₈	1.96 a	11.18 a	42.25 a	3.36 a	10.45 a
T ₉	1.4 bc	9.41bc	36.28 cd	2.93 bc	9.03 bc
T ₁₀	1.39 bc	9.60 bc	39.37 b	2.92 bc	9.04 bc
T ₁₁	1.49 bc	9.38 bc	38.11 bc	2.77 bc	8.53 bc
T ₁₂	1.55 b	10.70 ab	39.57 ab	2.83 bc	8.73 bc
LSD _(0.05)	0.1773	1.3169	2.8112	0.3329	1.0606
CV	7.07	8.13	4.54	6.85	7.06

Mean values within a column followed by the same letter do not differ significantly by Fisher's protected LSD test at $p \leq 0.05$

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]




Figure 4. Correlation graphs showing the reduced bulbing ratio in onion with relation to enhanced plant growth [plant height (A), number of leaves plant⁻¹ (B), number of roots plant⁻¹ (C) and root length (D)] recorded at 90 days after transplanting upon application of chitosan and plant probiotic bacteria

Clybio + *Paraburkholderia*, T₉: Chitosan @ 200 ppm, T₁₀: Chitosan @ 200 ppm + Clybio, T₁₁: Chitosan @ 200 ppm + *Paraburkholderia* and T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*. Required amount of chitosan was dissolved to 2% acetic acid and diluted in distilled water followed by heating and agitating for 30 minutes and rested to get normal temperature and pH being adjusted at 6.2 through adding 0.1N NaOH. Nutrient free *Paraburkholderia fungorum* BRRh-4 suspension at a concentration of approximately 1.5×10^9 CFU mL⁻¹ was prepared with modifying the method of Rahman *et al.* (2018b) where bacterial isolates were cultured in 1000 mL nutrient broth in conical flask and agitated at 80 rpm and 27 °C for 72 hours in incubator. The broth was centrifuged at 12,000 g and the pellet was washed thrice with sterilized distilled water to remove nutrients. Clybio made up of Yeast, *Lactobacillus* and *Bacillus natto* bacteria with the power of enzyme (indigenous microbial nutrient such as amino acid and peptide-based hormone substances) being collected from *Compass Corporation* was diluted to 0.1% concentration by adding 1 mL of clybio solute to 1000 mL of distilled water. The treatments were applied as foliar spray thrice at 20 days

intervals starting from 20 days after transplanting i.e., at 20 days, 40 days and 60 days after transplanting (DAT).

Experimental design and layout

The experiment was laid out in randomized complete block design (RCBD) with three replications. The experiment's field was first divided into 3 blocks, each block containing 12 unit plots represented a replication where each plot received a treatment and thereby a total of 36 unit plots measuring 1.8 m × 1.8 m were prepared where the 12 treatments were randomly assigned in the 12 plots of a block. The distance between the blocks was 50 cm and between the plots was 30 cm with the plant spacing of 15 cm × 15 cm to facilitate different intercultural operations. As many as 144 plants were accommodated in each unit plot and a total of 5184 plants were planted in the 36 unit plots.

Crop culturing and management

Prior to sowing the seeds were soaked overnight in water and allowed to sprout in a piece of cloth kept under the sunshade for two days. The sprouted seeds were sown directly on a raised seedbed in the nursery on 25th

Table 5. Effect of chitosan and plant probiotic bacteria on physiological weight loss (PLW) (%) of onion at 10 to 90 days after harvest

Treatment	Physiological loss in weight (%) at different days after harvest (DAH)								
	10DAH	20DAH	30DAH	40DAH	50DAH	60DAH	70DAH	80DAH	90DAH
T ₁	7.64 a	10.43 a	11.73 a	13.18 a	15.31 a	17.34 a	18.68 a	20.43 a	20.60 a
T ₂	3.43 c-f	5.44 cd	8.32 b	10.64 b	11.67 bc	12.27 bc	14.48 b	13.31 b	10.34 b
T ₃	4.02 b-d	5.64 b-d	8.17 b	9.53 bc	12.04 b	13.34 b	13.25 bc	12.22 bc	9.72 b
T ₄	4.11 bc	5.02 c-e	7.72 bc	9.22 b-d	10.49 b-d	11.55 c	12.51 c	10.41 cd	9.16 b
T ₅	2.49 f	3.31 f	3.88 e	5.48 g	7.15 g	9.15 de	8.48 e	6.60 fg	4.97 c
T ₆	2.79 ef	4.63 de	4.12 e	5.86 fg	7.42 g	9.33 de	10.65 d	9.20 de	6.28 c
T ₇	4.76 b	6.70 b	6.80 cd	8.63 c-e	10.06 c-f	11.84 c	12.57 c	10.91 de	9.26 b
T ₈	2.85 d-f	4.05 ef	4.67 e	5.73 fg	7.18 g	8.40 e	9.25 de	6.14 g	4.63 c
T ₉	3.74 b-e	5.70 b-d	6.39 d	7.28 ef	8.48 e-g	10.16 d	10.63 d	9.16 de	6.38 c
T ₁₀	3.90 b-e	5.27 cd	6.41 d	7.76 de	8.80 d-g	9.45 de	10.53 d	9.34 de	8.87 b
T ₁₁	3.70 b-e	5.77 bc	7.13 b-d	8.73 c-e	10.23 b-e	12.34 bc	12.68 bc	10.31 c-e	9.15 b
T ₁₂	3.54 c-f	5.06 c-e	6.22 d	7.70 de	8.31 fg	9.83 d	10.35 d	8.37 ef	6.25 c
LSD _(0.05)	1.1947	1.1213	1.2713	1.6165	1.8919	1.1063	1.8375	2.0194	1.8059
CV	18.02	11.86	11.05	11.49	11.45	5.81	9.04	11.32	12.12

Mean values within a column followed by the same letter do not differ significantly by Fisher's protected LSD test at $p \leq 0.05$. [T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]

Table 6. Effect of chitosan and plant probiotic bacteria on rotten bulbs (%) of onion at different days after harvest (DAH)

Treatment	Rotten bulbs (%)		
	30 DAH	60 DAH	90 DAH
T ₁	12.21 a	14.5 a	13.04 a
T ₂	7.18 c-e	10.68 cd	9.31 b
T ₃	7.11 c-e	10.62 c-e	7.78 c
T ₄	7.44 c-e	11.04 c	7.15 cd
T ₅	6.41 e	9.66 ef	5.75 f
T ₆	6.71 de	9.90 d-f	5.93 f
T ₇	9.94 b	12.14 b	7.43 cd
T ₈	6.28 e	9.38 f	5.66 f
T ₉	7.18 c-e	10.42 c-e	5.83 f
T ₁₀	8.76 bc	11.21 bc	6.93 de
T ₁₁	6.7 de	9.90 d-f	6.23 ef
T ₁₂	8.12 cd	11.2 bc	5.87 f
LSD _(0.05)	1.6613	0.9709	0.7076
CV	12.52	5.27	5.77

Mean values within a column followed by the same letter do not differ significantly by Fisher's protected LSD test at $p \leq 0.05$

[T₁: Control; T₂: Clybio; T₃: *Paraburkholderia*; T₄: Clybio + *Paraburkholderia*; T₅: Chitosan @ 100 ppm; T₆: Chitosan @ 100 ppm + Clybio; T₇: Chitosan @ 100 ppm + *Paraburkholderia*; T₈: Chitosan @ 100 ppm + Clybio + *Paraburkholderia*; T₉: Chitosan @ 200 ppm; T₁₀: Chitosan @ 200 ppm + Clybio; T₁₁: Chitosan @ 200 ppm + *Paraburkholderia*; T₁₂: Chitosan @ 200 ppm + Clybio + *Paraburkholderia*]




Figure 5. Correlation graphs showing the bulb yield enhancement of onion as a function of increased plant height (A), number of leaves/plant (B), number of roots/plant (C), root length (D) and reduced bulbing ratio (E) and shelf life extension with increased bulb weight (F) upon application of chitosan and plant probiotic bacteria




Figure 6. Average monthly maximum and minimum temperature, mean monthly relative humidity and monthly total rainfall of the growing area during November 2019 to June 2020 (Source: Department of Agricultural Engineering, BSMRAU)

November 2019. Upon germination (4-5 days after sowing), the young seedlings were exposed to dew by night and mild sunshine in the morning and evening. On 30th December 2019, 30-day old healthy, disease-free and uniform seedlings were transplanted to the main field with the spacing of 15 cm × 15 cm accommodating 144 plants in each unit plot. All intercultural operations like fertilizer management, insect-pest management, irrigation, weeding, mulching, etc were practiced as per guidelines and requirements. Before 7 days of harvest, when the plants attained maturity by showing drying up of leaves and weakening of necks, they were bent at the soil level by hands and kept as such up to harvest to fasten bulb maturity. Onion bulb was harvested by lifting with spade and hand on 25th April 2020. Care was taken so that no bulb was injured during lifting. Curing of onion was done for three days under a shed. Tops were separated from bulbs with a knife leaving 2 cm neck at ambient temperature (25.6 ± 2.6)⁰ C and then bulbs were stored in a well-ventilated house (25 ± 1⁰ C) separately as per treatments.

Measurement of growth, yield and storage qualities

Plant vegetative growth is an important indicator for reproductive performance. At the present study with onion, vegetative parameters like plant height (cm), number of leaves per plant, leaf length (cm), leaf width (mm), neck diameter (mm), number of roots, root length (cm) and bulb diameter (cm) were measured and recorded. Bulbing ratio was also calculated through dividing the neck diameter by bulb diameter. The growth parameters were measured from randomly selected five onion plants of each treatment plot and averages were used. All the data were gathered at 30, 60 and 90 days after transplanting (DAT). Again, after harvest total number of bulbs, individual bulb weight (g), yield per plot (kg) and yield per hectare (t) were estimated by a weighing balance. Splitting of bulb (%) was recorded by the following formula-

$$\text{Splitting of bulb (\%)} = \frac{\text{Number of split bulbs} \times 100}{\text{Number of total bulbs (plants)}}$$

Physiological loss in weight (PLW) (%) and rotten bulb (%) were also recorded. The following formula were used for calculating PLW (%) and rotten bulb (%) –

$$\text{PLW (\%)} = \frac{\text{Fresh or initial weight} - \text{Storage or final weight}}{\text{Initial weight}} \times 100$$

$$\text{Rotten bulb (\%)} = \frac{\text{Number of rotten bulbs} \times 100}{\text{Total number of bulbs}}$$

Shelf life of onion in storage was determined by keeping the bulbs in ambient temperature ($25 \pm 1^\circ \text{C}$) in normal storage conditions at the laboratory and observing the bulb quality up to optimum marketing.

Statistical analysis

The collected data were tabulated and analyzed statistically using MSTAT-C statistical package program. The treatment means were separated by Fisher's Least Significant Difference (LSD) test, using a p-value of ≤ 0.05 to be statistically significant (Gomez and Gomez, 1984). Correlation analysis with the treatment means were also performed for bulbing ratio, bulb yield and shelf life in relation to different vegetative growth parameters observed at 90 days after transplanting using MS Office Excel 2007.

Conclusion

Utilization of naturally occurring chitosan biopolymer and plant probiotic bacteria has been practiced for quality enhancement of produce with multiple advantages to human health and environment. The present investigation exhibited that shrimp shell chitosan, clybio and *Paraburkholderia* in combinations enhance the vegetative and reproductive growth of onion. Among the combinations, 3-time foliar spray Chitosan (100 ppm) + Clybio (0.1%) + *Paraburkholderia* ($1.5 \times 10^9 \text{ CFU mL}^{-1}$) significantly enhanced the growth as well as bulb production of onion at field level under semi-arid condition in Bangladesh. Furthermore, the same treatment also responded for maximum shelf life through minimizing physiological weight loss and protecting rotting of onion bulb in storage.

Acknowledgements

The authors would like to express their heartiest gratitude to the Department of Horticulture, Bangabandhu Sheikh Mujibur Rahman Agricultural University (BSMRAU), Gazipur, Bangladesh for the giving opportunity to accomplish the whole research activities; Institute of Biotechnology and Genetic Engineering, BSMRAU for offering some of the laboratory facilities and Bangladesh Agricultural Research Institute, Gazipur, Bangladesh for supplying with the planting materials.

References

Abdel-Mawgoud AM, Tantawy AS, El-Nemr MA, Sassine YN (2010) Growth and yield responses of strawberry plants to chitosan application. *Eur J Sci Res.* 39(1): 170-177.

Abdel-Moneim MM, El-Mazny MY, Abdel-Mageed YT, Moustafa YMM, Yamani SHS (2015) Effect of some natural antioxidants on the productivity and storage ability of Egyptian onion grown in sandy soil. In *Book of Proceedings 2nd International Conference for Agriculture and Irrigation in the Nile Basin Countries* (pp. 23-25).

Afify A, Hauka F, Elsawah A (2018) Plant growth-promoting rhizobacteria enhance onion (*Allium cepa* L.) productivity and minimize requisite chemical fertilization. *Env Biodivers Soil Secur.* 2(2018): 119-129.

Ahmed MEM, Farm E (2015) Response of garlic plants (*Allium sativum* L.) to foliar application of some biostimulants. *Egypt J Hort.* 42(1): 613-625.

Akter S, Uddin AFMJ, Hossin I, Islam MN (2021) Influence of seed priming and clybio application on growth and yield of spinach. *Int J Agric Environ Bio-res.* 6(3): 223-227.

Axel C, Zannini E, Coffey A, Guo J, Waters DM, Arendt EK (2012) Ecofriendly control of potato late blight causative agent and the potential role of lactic acid bacteria: a review. *Appl Microbiol Biotechnol.* 96: 37-48.

Bashan Y, de-Bashan LE (2005) Bacteria/plant growth-promotion. In: Hillel D (ed). *Encyclopaedia of Soils in the Environment*. Oxford: Elsevier, pp. 103–115.

Basit A, Hassnain MA, Ullah I, Shah ST, Zuhair SA, Ullah I (2020) Quality indices of tomato plant as affected by water stress conditions and chitosan application. *Pure Appl Biol.* 9(2): 1364-1375.

Chakraborty M, Islam T (2022) 12-Growth-stimulating effects of chitosan biopolymer in plants. In *Biostimulants for Crop Production and Sustainable Agriculture*. 179.

Colo JOSIP, Hajnal-Jafari T, ĐURIĆ S, Stamenov D, Hamidović S (2014) Plant growth promotion rhizobacteria in onion production. *Pol J Microbiol.* 63(1): 83.

Dilpreet T, Kulbir S, Varinder S, Jagdish S (2017) Growth, yield and quality of onion as influenced by integrated nutrient management. *Int J Agric Sci Res.* ISSN 0975-3710.

El-Miniawy SM, Ragab ME, Youssef SM, Metwally AA (2013) Response of strawberry plants to foliar spraying of chitosan. *Res J Agric Biol Sci.* 9(6): 366-372.

Emami Bistgani Z, Siadat SA, Bakhshandeh A, Ghasemi Pirbalouti A, Hashemi M (2017) Morpho-physiological and phytochemical traits of (*Thymus daenensis* Celak.) in response to deficit irrigation and chitosan application. *Acta Physiol Plant.* 39: 1-3.

FAO (1999) *FAO quarterly bulletin of statistics*. Food and Agricultural Organization of the United Nations, Rome, Italy. 12: 91.

Fawzy ZF, El-Magd AM, Li Y, Ouyang Z, Hoda AM (2012) Influence of foliar application by EM" effective microorganisms", amino acids and yeast on growth, yield and quality of two cultivars of onion plants under newly reclaimed soil. *J Agric Sci* 4(11): 26.

Garcia-Secco D, Zhang Y, Gutierrez-Manero FJ, Martin C, Ramos-Solano B (2015) Application of *Pseudomonas fluorescens* to blackberry under field conditions improves fruit quality by modifying flavonoid metabolism. *PLoS One.* 10(11):e0142639.

Geries LSM, Omnia SM, Hashemb, Marey RA (2020) Soaking and foliar application with chitosan and nano chitosan to enhancing growth, productivity and quality of onion crop. *Plant Arch.* 20(2): 3584-3591.

Giassi V, Kiritani C, Kupper KC (2016) Bacteria as growth-promoting agents for citrus rootstocks. *Microbiol Res.* 190: 46-54.

Gomez KA, Gomez AA (1984) *Statistical Procedures for Agricultural Research*. John Wiley and Sons, Inc., New York. 67-215.

Hamed HA, Moustafa YA, Abdel-Aziz SM (2011) In vivo efficacy of lactic acid bacteria in biological control against *Fusarium oxysporum* for protection of tomato plant. *Life Sci J.* 8(4): 462-468.

- Hassnain M, Alam I, Ahmad A, Basit I, Ullah N, Alam I, Ullah MA, Khalid B, Shair MM. (2020) Efficacy of chitosan on performance of tomato (*Lycopersicon esculentum* L.) plant under water stress condition. *Pak J Agric Res.* 33(1):27-41.
- Ikedda DM, Weinert E, Chang KC, McGinn JM, Miller SA, Kelihoomalulu C, DuPonte MW (2013) Natural farming: lactic acid bacteria. *Sustain Agric.* 8: 3-4.
- Islam M, Hossain M (2012) Plant probiotics in phosphorus nutrition in crops, with special reference to rice. *Bacteria in agrobiology: plant probiotics.* pp. 325-363.
- Jiménez-Gómez A, Celador-Lera L, Fradejas-Bayón M, Rivas R (2017) Plant probiotic bacteria enhance the quality of fruit and horticultural crops. *AIMS Microbiol.* 3(3): 483.
- Jiménez-Gómez A, Menéndez E, Flores-Félix JD, García-Fraile P, Mateos PF, Rivas R (2016) Effective colonization of spinach root surface by *Rhizobium*. In *Biological Nitrogen Fixation and Beneficial Plant-Microbe Interaction*, pp. 109-122, Springer International Publishing.
- Jitareerat P, Paumchai S, Kanlayanarat S, Sangchote S (2007) Effect of chitosan on ripening, enzymatic activity, and disease development in mango (*Mangifera indica*) fruit. *New Zealand J Crop Hortic.* 35(2): 211-218.
- Karlidag H, Esitken A, Yildirim E, Donmez MF, Turan M (2010) Effects of plant growth promoting bacteria on yield, growth, leaf water content, membrane permeability, and ionic composition of strawberry under saline conditions. *J Plant Nutr.* 34(1): 34-45.
- Kloepper JW, Ryu CM, Zhang S (2004) Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathol.* 94(11): 1259-1266.
- Kuete V (ed) (2017) *Allium cepa, Medicinal spices and vegetables from Africa: therapeutic potential against metabolic, inflammatory, infectious and systemic diseases.* Academic Press. 353-361.
- Kumar A, Ram RB, Maji S, Kishor S, Yadav R, Meena KR (2017) Effect of organic manures, biofertilizers and micronutrients on growth, yield and quality of onion (*Allium cepa* L.). *Int J Agric Sci.* 13(2): 236-241.
- Liu Y, Wisniewski M, Kennedy JF, Jiang Y, Tang J, Liu J (2016) Chitosan and oligochitosan enhance ginger (*Zingiber officinale* Roscoe) resistance to rhizome rot caused by *Fusarium oxysporum* in storage. *Carbohydr Polym.* 151: 474-479.
- Mahmood N, Abbasi NA, Hafiz IA, Ali I, Zakia S (2017) Effect of biostimulants on growth, yield and quality of bell pepper cv. *Yolo Wonder*. *Pak J Agric Sci.* 54(2).
- Marrelli M, Amodeo V, Statti G, Conforti F (2019) Biological properties and bioactive components of *Allium cepa* L.: Focus on potential benefits in the treatment of obesity and related comorbidities. *Molecules.* 24(1): 119.
- Matter FMA, Abou-Sreea AIB (2016) Influence of application methods of bio-fertilization on vegetative growth, seed yield and chemical composition of Fenugreek plants. *Egypt. J Hort.* 43(1): 19-33.
- Mehebab MS, Mahmud NU, Rahman M, Surovy MZ, Gupta DR, Hasanuzzaman M, Islam MT (2019) Chitosan biopolymer improves the fruit quality of litchi (*Litchi chinensis* Sonn.). *Acta Agrobot.* 72(2): 1773.
- Mohite B (2013) Isolation and characterization of indole acetic acid (IAA) producing bacteria from rhizospheric soil and its effect on plant growth. *J Soil Sci Plant Nutr.* 13(3): 638-49.
- Mondal MMA, Malek MA, Puteh AB, Ismail MR, Ashrafuzzaman M, Naher L (2012) Effect of foliar application of chitosan on growth and yield in okra. *Aust J Crop Sci.* 6(5): 918-921.
- Mukta JA, Rahman M, Sabir AA, Gupta DR, Surovy MZ, Rahman M, Islam MT (2017) Chitosan and plant probiotics application enhance growth and yield of strawberry. *Biocatal Agric Biotechnol.* 11: 9-18.
- Parr JF, Hornick SB, Whitman CE (1991) First International Conference on Kyusei Nature Farming. Proceedings of the Conference held October 17-21, 1989, at Khon Kaen, Thailand. 175 p.
- Parvin MA, Zakir HM, Sultana N, Kafi A, Seal HP (2019) Effects of different application methods of chitosan on growth, yield and quality of tomato (*Lycopersicon esculentum* Mill.). *Arch Agri Sci J.* 4(3): 261-267.
- Rahman M, Mukta JA, Sabir AA, Gupta DR, Mohi-Ud-Din M, Hasanuzzaman M, Miah MG, Rahman M, Islam MT (2018a) Chitosan biopolymer promotes yield and stimulates accumulation of antioxidants in strawberry fruit. *PLoS One.* 13(9): p.e0203769.
- Rahman M, Sabir AA, Mukta JA, Khan M, Alam M, Mohi-Ud-Din M, Miah M, Rahman M, Islam MT (2018b) Plant probiotic bacteria *Bacillus* and *Paraburkholderia* improve growth, yield and content of antioxidants in strawberry fruit. *Sci Rep.* 8(1): 1-11.
- Rashid MHA, Massiah AJ, Thomas B (2015) Genetic regulation of day length adaptation and bulb formation in onion (*Allium cepa* L.). In VII International Symposium on Edible Alliaceae. 1143: 7-14.
- Reetha S, Bhuvaneshwari G, Thamizhiniyan P, Mycin TR (2014) Isolation of indole acetic acid (IAA) producing rhizobacteria of *Pseudomonas fluorescens* and *Bacillus subtilis* and enhance growth of onion (*Allium cepa* L.). *Int J Curr Microbiol Appl Sci.* 3(2): 568-574.
- Rodríguez AF, Costales D, Peña DG, Morales D, Mederos Y, Jerez E, Pino JC (2017) Chitosans of different molecular weight enhance potato (*Solanum tuberosum* L.) yield in a field trial. *Span J Agric Res.* 15(1): 25.
- Romanazzi G, Feliziani E, Baños SB, Sivakumar D (2017) Shelf life extension of fresh fruit and vegetables by chitosan treatment. *Crit Rev Food Sci Nutr.* 57(3): 579-601.
- Ruzzi M, Aroca R (2015) Plant growth-promoting rhizobacteria act as biostimulants in horticulture. *Sci. Hortic.* 196: 124-134.
- Shaheen AM, Ragab ME, Rizk FA, Mahmoud SH, Soliman MM, Omar NM (2019) Effect of some active stimulants on plant growth, tubers yield and nutritional values of potato plants grown in newly reclaimed soil. *J Anim Plant Sci.* 29(1): 215-225.
- Shehata SA, Fawzy ZF, El-Ramady HR (2012) Response of cucumber plants to foliar application of chitosan and yeast under greenhouse conditions. *Aust J Basic Appl Sci.* 6(4): 63-71.
- Shrestha A, Kim BS, Park DH (2014). Biological control of bacterial spot disease and plant growth-promoting effects of lactic acid bacteria on pepper. *Biocontrol Sci Technol.* 24(7): 763-79.
- Soppelsa S, Kelderer M, Casera C, Bassi M, Robatscher P, Matteazzi A, Andreotti C (2019). Foliar applications of biostimulants promote growth, yield and fruit quality of strawberry plants grown under nutrient limitation. *Agronomy.* 9(9): 483.
- Steer BT (1980) The bulbing response to day length and temperature of some Australasian cultivars of onion (*Allium cepa* L.). *Aust J Agric Res.* 31(3): 511-518.

- Sultana S, Islam M, Khatun MA, Hassain M, Huque R (2017) Effect of foliar application of oligo-chitosan on growth, yield and quality of tomato and eggplant. *Asian J Agric Res.* 11(2): 36-42.
- Sutariati GAK, Khaeruni A, Madiki A, Rakian TC, Mudi L, Fadillah N (2019) Seed biopriming with indigenous endophytic bacteria isolated from Wakatobi rocky soil to promote the growth of onion (*Allium ascalonicum* L.). In *IOP Conference Series: Earth and Environmental Science* (Vol. 260, No. 1, p. 012144). IOP Publishing.
- Tinna D, Garg N, Sharma S, Pandove G, Chawla N (2020) Utilization of plant growth promoting rhizobacteria as root dipping of seedlings for improving bulb yield and curtailing mineral fertilizer use in onion under field conditions. *Sci Hortic.* 270: 109432.
- Tsavkelova EA, Klimova SY, Cherdyntseva TA, Netrusov AI (2006) Microbial producers of plant growth stimulators and their practical use: a review. *Appl Biochem Microbiol.* 42(2): 117-126.
- Uddin AFMJ, Nasif FM, Kaynat B, Maliha M, Rakibuzzaman M (2021) Organic Bio-stimulator Application on Growth and Yield of Strawberry. *Int J Acad Res Bus Soc Sci.* 9(3): 01-05.
- Uthairatanakij A, Teixeira da Silva JA, Obsuwan K (2007) Chitosan for improving orchid production and quality. *Orchid Sci Biotechnol.* 1(1): 1-5.
- Ware M (2017) <https://www.medicalnewstoday.com/articles/276714.php> . (2019).
- Yildirim E, Turan M, Ekinci M, Dursun A, Gunes A, Donmez M (2015) Growth and mineral content of cabbage seedlings in response to nitrogen fixing rhizobacteria treatment. *Rom Biotech Lett.* 20: 10929-10935.