Australian Journal of Crop Science

AJCS 16(01):128-136 (2022) doi: 10.21475/ajcs.22.16.01.p3372



Understanding the flux of nitrous oxide from the eucalypt soil in monoculture and Integrated Crop–Livestock–Forest systems

Júlia Graziela da Silveira¹, Charlote Wink², Fernando Vieira Cesário³, Natassia Magalhães Armacolo⁴, Anna Karolyne da Silva Nogueira⁵, Jacqueline Jesus Nogueira da Silva⁶, Fernanda Figueiredo Granja Dorilêo Leite⁷, Marcela Cardoso Guilles da Conceição⁶, Silvio Nolasco de Oliveira Neto¹, Renato de Aragão Ribeiro Rodrigues^{7,8*}

¹Departamento de Engenharia Florestal, Universidade Federal de Viçosa, Viçosa, 36570-977, Minas Gerais, Brasil ²Instituto de Ciências Agrárias e Ambientais, Universidade Federal de Mato Grosso, Sinop, 78557-267, Mato Grosso, Brasil

³Doutor em Geografia

⁴Centro de Ciências Agrárias, Universidade Estadual de Londrina, Londrina, 86057-970, Paraná, Brasil

⁵Mestra em Agronomia

⁶Doutora em Geociências

⁷Departamento de Geoquímica, Universidade Federal Fluminense, Niterói, 24020-140, Rio de Janeiro, Brasil

⁸Embrapa Solos, Jardim Botânico, Rio de Janeiro, 22460-000, Rio de Janeiro, Brasil.

*Corresponding author: renato.rodrigues@embrapa.br

Abstract

The Brazilian agricultural sector is largely responsible for nitrous oxide (N_2O) soil emissions, mainly due to beef cattle and the use of synthetic nitrogen fertilizers. Therefore, Brazil is looking for measures, such as integrated crop–livestock– forest (ICLF), to increase productivity and reduce greenhouse gas emissions in this sector. The forest component within this system plays a positive role in the context of climate change, soil conservation, carbon dioxide (CO_2) sequestration, and biodiversity protection. The aim of this study was to evaluate the effect of management and rainfall on N_2O emissions in eucalypt monoculture soils and eucalypt soils in ICLF systems. Manual static chambers were used to collect gas samples, from November 2013 to October 2014, in four treatments, i.e., one eucalypt monoculture (F) and three modalities of ICLF (livestock–forest [LF], livestock–crop–forest [LCF], and integrated crop–livestock–forest [ICLF]). A gas chromatograph with an electron capture detector was used to measure the N_2O concentrations. The results showed that rainfall considerably affected N_2O fluxes across all the treatments, indicating that rainfall is the main factor in increasing emissions. During the wet season, the N_2O levels ranged from 0.158 to 0.482 kg $N-N_2O$ ha⁻¹ across all treatments. During the dry season, all treatments behaved like sinks of N_2O . Moreover, N_2O flux did not differ between the soils in the eucalypt monoculture and ICLF systems. This indicates that the forestry component in the ICLF systems did not affect N_2O soil fluxes.

Keywords: agroforestry system, greenhouse gases, integrated systems, mitigation, sustainability.

Abbreviations: GHG_greenhouse gases; CO_2 _carbon dioxide; N_2O_n itrous oxide; CH4_methane; N_nitrogen; NDC_Nationally Determined Contribution; CO_2eq_Carbon dioxide equivalent; ICLF_integrated crop-livestock-forest; NO^{3-}_N itrate; LF_livestock-forest; LCF_livestock-crop-forest; F_eucalypt monoculture; WFPS_water filled pore spaces; N_2 _elementar nitrogen.

Introduction

Atmospheric concentrations of the three main greenhouse gases (GHG), carbon dioxide (CO₂), nitrous oxide (N₂O), and methane (CH₄), increased by 146%, 257%, and 122%, respectively, compared to pre-industrial levels (WMO, 2018) due to land use change and agricultural practices, among other causes (Muller, 2005; Hartmann et al., 2013; Rodrigues et al., 2017; Tian et al., 2020).

In Brazil, the agricultural sector has the largest share (33.6%) of the net GHG emissions in 2016, reaching up to 439.213 Gg CO_2eq , showing an increase of 2.3% compared to that in 2015 (Brasil, 2020). In terms of the regions associated with

high emissions, the Midwest and Mato Grosso State are leading, accounting for 34% and 12% of the total emissions, respectively, with the main sources of emissions being beef cattle and the use of synthetic nitrogen fertilizers (Silveira et al., 2018).

Although atmospheric concentrations of all three gases are relevant in agriculture, major emphasis should be placed on N₂O, as its global warming potential is 265 times higher than that of CO₂ (Myhre et al., 2013). N₂O emissions tend to increase with the expansion of agricultural sector emerging economies, such as Brazil, where the agriculture sector is the

largest contributor to GHG emissions (Brasil, 2020; Tian et al., 2020). This gas is the result of soil nitrification and denitrification processes, which vary with the dynamics of nitrogen (N) (Saggar, 2010; Ussiri and Lal, 2013) and is also emitted due to the decomposition of plant residues or application of nitrogen fertilizers (Sato et al., 2017; Scott et al., 2018). Other agronomic management practices (e.g., irrigation and the use of manure) (Reay et al., 2012; Bayer et al., 2015; Tian et al., 2015; Martins et al., 2015), as well as cultivated crops and local rainfall regimes may also add to the affect (Li et al., 2016).

To increase efforts to mitigate climate change, Brazil, through its Nationally Determined Contributions (NDCs), which include mitigation, adaptation, and implementation components, has committed to reducing GHG emissions by 37% by 2025 and 43% by 2030, below 2005 levels (BRASIL, 2015). By 2030, actions under the NDCs are anticipated to expand the area under integrated crop–livestock–forest (ICLF) system by 5 million hectares (BRASIL, 2015). Partial results of these public policies are already obvious, as according to the ICLF Network, Brazil went from having an area of 1.87 million hectares under ICLF systems in 2005 to 11.5 million hectares by the 2015/2016 harvest, with the potential to stock 35.1 million tons of CO_2eq (Rede ILPF, 2016).

The ICLF system has been identified as an efficient land management strategy that provides environmental, economic, and social benefits, such as soil and water conservation, wood production, and improved animal welfare, which can increase milk and beef production (Cordeiro et al., 2015; Alves et al., 2017). Additionally, it can improve the restoration of degraded pastures, mitigate GHG emissions, and increase soil carbon (C) sequestration by increasing biomass above and below the ground (Cerri et al., 2007; Euclides et al., 2010; Nair, 2012; Dube et al., 2012; Paula et al., 2013; Carvalho et al., 2014; Salton et al., 2014; Paustian et al., 2016; Stocker et al., 2013; Zomer et al., 2016). Furthermore, the use of crop rotations in ICLF with variable nitrogen availability is also a strategy for mitigating N₂O emissions (Benoit et al., 2015; Jain et al., 2016).

According to Figueiredo et al. (2017), ICLF can reduce GHG emissions mainly due to pasture improvements, increase in the livestock yield, and the potential for carbon sinks in soil and biomass to offset emissions related to livestock management. In addition, these authors report that emissions can be reduced in terms of CO2eq emitted per kg of bovine weight produced, increasing meat, grain, and wood production. According to Nogueira et al. (2016), the cumulative N₂O emissions in ICLF (0.4 kg N ha⁻¹) may be lower than that in a monoculture crop $(1.4 \text{ kg N} \text{ ha}^{-1})$, and similar to pasture monoculture, when there is no livestock $(0.35 \text{ kg N ha}^{-1})$, thereby presenting a potential to mitigate N₂O emissions. Moreover, the adoption of ICLF can help achieve both environmental protection and the development of more efficient and sustainable agriculture in states with highly intensive livestock systems, low stocking rates, and where the agricultural sector is expanding rapidly, such as the Mato Grosso State (the country's leading cattle and soy producer) (Gil et al., 2015).

The lower N_2O emission from forest areas compared to the crop areas is related to the greater ability of the tree roots to absorb water and NO^{3-} in depth, reducing denitrification and leaching (Amadi et al., 2016). The nitrogen absorbed by the tree roots is returned to the soil through the leaf litter, resulting in more efficient N cycling and thus, decreasing the demand for N and emission of N₂O (Thevathasan et al.,

2012). Given the advantages of the ICLF, the forest component deserves attention as it plays a positive role in control of climate change and deforestation through restoration of degraded areas, soil conservation, CO_2 sequestration, biodiversity protection, and reduction of pressure for the use of native forests in industrial processes (Cuer et al., 2018).

Of the total planted forest area in Brazil, 5.7 million hectares are eucalypt forests (Cuer et al., 2018), which indicates the importance of these species for the sector. These are fastgrowing trees with high carbon sequestration potential during development (Burrows et al., 2002; Du et al., 2015; Bauters et al., 2019). Eucalypt plantations have been reported as a source of N₂O and CO₂ and a sink for CH₄ in semi-arid and subtropical climates, as observed in most forest ecosystems (Zhang et al., 2017).

However, GHG fluxes in eucalypt plantations have not been well described in ICLF systems. Thus, in the present study, two questions were posed:

1. Do the fluxes of soil N_2O differ between eucalypt monocultures and eucalypt in ICLF systems?

2. Does rainfall affect N₂O emissions in a similar manner in both eucalypt monocultures and ICLF systems?

The implementation of an integrated management system is expected to reduce N_2O emissions due to the increased inclusion of different crops, favoring increased productivity and efficiency in nutrient utilization. Rainfall is also expected to have a positive effect on N_2O emissions owing to the filling of porous soil spaces and the stimulation of microbial action, thereby altering N_2O production.

Results

The N_2O fluxes from eucalypt soil in the monoculture and in the integrated systems had the same variation throughout the year, as shown in Figure 1. The months from October to May showed positive N_2O fluxes, while the months from June to September showed negative fluxes across all systems. In this study, the data show an evident seasonal effect of rainfall on N_2O fluxes.

The monthly variation of N₂O fluxes throughout the year in the study areas did not exceed 0.10 kg N-N₂O month⁻¹. In the dry season (from May to September), when soils have low humidity due to low rainfall, all systems acted as N₂O sinks, except in May for ICLF, livestock–forest (LF), and livestock– crop–forest (LCF) (Figure 1). The cumulative highest average monthly N₂O flux observed during the period was approximately 0.08 kg N-N₂O month⁻¹ for the eucalypt soil in the ICLF system in May, while the highest consumption potential was -0.05 kg N-N₂O month⁻¹ for the eucalypt soil in the LF system during June (Figure 1).

There was a significant difference between the cumulative emissions of the dry and wet seasons for all the treatments (p = 0.0000415) (Figure 2a). Overall, the dry season shows negative emission values, indicating that the soil acts as a sink for N₂O. Furthermore, there were no significant differences between the treatments in the dry season (p = 0.1129), varying from -0.022 (ICLF) to -0.156 kg N-N₂O ha⁻¹ (LF). Similarly, in the wet season, there was no significant difference between the treatments (p = 0.0987), and the emission varied from 0.158 (LCF) to 0.482 kg N-N₂O ha⁻¹ (F) (Figure 2a).

The results from analyses of the net cumulative emissions during the entire period shows that eucalypt soil from all the



Figure 1. Cumulative average monthly N-N₂O fluxes (lines), and standard deviation (whiskers), and rainfall (bars) throughout the year for each treatment. F_eucalypt monoculture; ICLF_integrated crop-livestock-forest; LCF_livestock-crop-forest; LF_livestock-forest.



Figure 2. Cumulative $N-N_2O$ soil emissions between dry and wet seasons for each treatment (a) and net cumulative emission between the wet and dry period for each treatment (b). The vertical bars represent the standard deviation. F_eucalypt monoculture; ICLF_integrated crop-livestock-forest; LCF_livestock-crop-forest; LF_livestock-forest.



Figure 3: Location of the study site in Mato Grosso, Brazil.



Figure 4. Aerial image of the four treatments and three replications evaluated in Sinop, Mato Grosso, Brazil. Source: Google Earth. ICLF_integrated crop-livestock-forest; LF_livestock-forest; LF_livestock-crop-forest; F_eucalypt monoculture.

			F	LF	LCF	ICLF		
						50.0 Kg ha ⁻¹ Soybean (<i>Glycine</i>		
چې	Sowing	Oct - 2011				Max L.) BRSGU 8560 RR	-	
				Hybrid <i>F</i>	= urophylla x F a	randis		
		Nov - 2011	200 kg ha ⁻¹ NPK (00-20-20)					
			350 kg ha ⁻¹ (Phosphate fertilizer)					
				400 Kg ha ⁻ ' N	JPK (04-30-16)			
				Brachiaria briza	ntha cv. Marandu			
			(6 kg ha ⁻ ')					
R	Ferti	Dec - 2011	96.40 Kg ha ⁻¹ NPK (20-00-20)				2011 - 2	
(@)	lization	Jan - 2012						
	Harvest					Soybean (John Deere STS 9470)	2012 Harvest	
چ ت	Sowing	Feb - 2012				Brachiaria brizantha cv. Marandu (5.50 kg ha ⁻¹)		
						10 Kg ha ⁻¹ Corn (<i>Zea may</i> s L.)		
						375 Kg ha ⁻¹ NPK (04-30-16)		
						300 kg ha ⁻¹ Urea		
Ø	Fertilization	Mar - 2012	20 Kg ha ⁻¹ NPK (20-00-20)		20 Kg ha ⁻¹ NPK (20-00-20)			
بينين ا	Harvest	Jul - 2012				Corn (John Deere STS 9470)		
۵	S					25 Kg ha ⁻¹ Soybean (<i>Glycine max</i>		
So and a second	owing	Oct - 2012				L.) BRSGO 8560 RR		
	E.					350 Kg Ha INPK (00-20-20)		
Ø	ertilization	Jan - 2013	119.0 Kg ha ⁻¹ NPK (20-00-20)					
₩ 	Harvest	Feb - 2013				Soybean (John Deere STS 9470)	2012 – 2013 Harves	
ي چ	Sowing					30 Kg ha ⁻¹ Corn (Zea mays L.)		
						5.50 kg ha ⁻¹ Seeds <i>Brachiaria</i>		
						300 Kg ha ⁻¹ NPK (08-28-16)		
~	Fe		96.0 Kg ba ⁻¹ NPK (20-00-20)			7		
	rtiliz	Mar- 2013	30.0 Kg na Ni K (20-00-20)					
	ation					210 Kg ha ⁻¹ (Urea)		
	Harvest	Jul - 2013				Corn (John Deere STS 9470)		
چ ا	Sow	Oct - 2013			25 Kg ha ⁻¹ So	ybean (<i>Glycine max</i> L.) BRSGO 8560RR		
	ving				Li	Liming (1500 kg ha ⁻¹)		
	т				400 kg ha ⁻¹ NPK (00-20-20) Soybean (John Deere STS 9470)		2013 – 2(
	larvest							
چ ج	Sowing	Feb - 2014			13.72 kg ha ⁻¹ Corn (<i>Zea mays</i> L.) DKB 390 Pro		014	
					6 kg ha ⁻¹ Seeds <i>Brachiaria brizantha</i> cv. Marandú		Harv	
					450 4	450 Kg ha ⁻¹ NPK (04-30-16)		
		Mar - 2014			300 Kg ha ⁻¹ (Ureia)			
	Harvest	Jul - 2014			Corn (John Deere STS 9470)			
چ	Sowing	Oct - 2014			25 Kg ha ⁻¹ So	ybean (<i>Glycine max</i> L.) BRSGO 560RR	2014 – 2015 Harvest	
					400 1	⟨g ha ⁻¹ NPK (00-20-20)		

Figure 5. Timeline and description of all management (i.e., Fertilizing, Sowing, Harvest) applied in the treatments, including all species planted and rotation used. The crop timeline for this study starts in the year 2011 (light gray) and finishes in 2015 (dark gray). F_eucalypt monoculture; LF_livestock-forest; LCF_livestock-crop-forest; ICLF_integrated crop-livestock-forest.

treatments had positive emission values, with no significant differences in the treatments (p = 0.1473) (Figure 2b). The emissions varied from 0.057 (LCF) to 0.336 kg N-N₂O ha⁻¹ (F).

Discussion

The predominantly positive N₂O fluxes observed during rainy season across all treatments may be a reflection of the availability of inorganic N due to fertilizer application and mineralization of plant residues (Bayer et al., 2015; Piva et al., 2019) from the other components of the integrated systems. In addition, the anaerobic conditions of the highly humid soil can directly influence the soil's water-filled pore spaces (WFPS), which are important in the denitrification processes and are the main factor in the formation of N₂O (Rosenkranz et al., 2006; Konda et al., 2008; Fang et al., 2012; Butterbach-Bahl et al., 2013). These processes may be related to the higher activity of microorganisms in soaked soils, potentiated when saturation level of WFPS is more than 70% (Weerden et al., 2012; Butterbach-Bahl et al., 2013; Corrêa et al., 2016). However, when the WFPS saturation is equal to or greater than 80%, the denitrification process is intensified, and the end product under these conditions will be mainly N₂ (and not N₂O) (Linn and Doran, 1984; Davidson et al., 2000; Bateman and Baggs, 2005; Butterbach-Bahl et al., 2013). Thus, the reduction in emissions after December (Figure 1) in ICLF, LCF, and LF may be related to high WFPS saturation due to high rainfall. Changes in soil moisture and temperature maybe responsible for approximately 95% of the temporal variations in field N₂O emissions (Bouwman, 1998). Nevertheless, higher N₂O consumption potential during periods of low rainfall may be associated with the low N content available in the soil, reducing the nitrification and denitrification activities, which are the main processes for N₂O emission (Chapuis-Lardy et al, 2007). Furthermore, low mineral nitrogen and rapid soil water drainage may not offer favorable conditions for high N₂O fluxes (Neves, 2016; Carvalho et al., 2017). Under these circumstances, air can diffuse into the macro- and micro-pores of the soil and allow microorganisms to use N₂O as a source of N (Rosenkranz et al., 2006). In their study evaluating GHG emissions with different N rates in eucalypt areas in China, Zhang et al. (2017) found higher emissions of nitrous oxide in the wet season and lower emissions in the dry season, with the highest emissions always observed for the treatments with higher N rates, even in the dry season. Thus, the low N₂O fluxes found in this study during the dry season may be related to the high soil porosity, drought conditions, and low soil N content. Our results showed a high standard deviation (Figure 2), as soil N₂O fluxes normally have high spatial and temporal variability due to the heterogeneity of environmental parameters and soil characteristics (McDaniel et al., 2017; Rivera et al., 2019; Charteris et al., 2020). The climate seasons also impact the N₂O emission, as found by Nascimento and Rodrigues (2019) who found positive cumulative fluxes in the dry period with approximately 0.25 kg N-N₂O ha⁻¹ and 0.50 kg N-N₂O ha⁻¹ in soils with eucalypt monoculture and in native forests, respectively. Their results differ from the results of this study, in which we showed that the eucalypt soil from all the treatments acted as a sink, with negative values (N₂O consumption) in the dry period (Figure 2a). In the study by Nascimento and Rodrigues (2019), the maximum cumulative value during the rainy season was 0.45 kg N-N₂O ha⁻¹ for the eucalypt monoculture and 0.50 kg $N-N_2O$ ha⁻¹ for the native forest. A comparison

between the above results and the results of this study showed that the soil of a eucalypt forest planted in an integrated system has a lower potential N_2O emission than that of a eucalypt monoculture (F).

The total N₂O emission for the entire period for the eucalypt soil is still low (Figure 2b) when compared to the total values of the native Amazon Forest soil (0.6 kg N-N₂O ha⁻¹) (Nascimento et al., 2020), native Atlantic Forest soil (0.82 kg N-N₂O ha⁻¹), and soil of eucalypt monoculture (0.55 kg N-N₂O ha⁻¹) (Silva, 2019). The results in this study are also moderate as compared to a study carried out in the transition biome Cerrado and Amazon Forest, which found cumulative values of approximately 1.0 kg N-N₂O ha⁻¹ for native forest and 0.70 kg N-N₂O ha⁻¹ for monoculture of eucalypt (Nascimento and Rodrigues, 2019).

Although we did not evaluate all the components of the integrated systems (crops, livestock, and forest), Nogueira et al. (2016) evaluated it in the same study area and found cumulative values of 0.4 kg N-N₂O ha⁻¹ for ICLF and 1.4 kg N-N₂O ha⁻¹ for monoculture crops, demonstrating the trend of low emissions in the eucalypt soil of integrated systems. In this study, animal non-grazing in the areas near eucalypt may also have been reflected in the low emissions, as there was no deposition of urine residues and cattle feces, which can influence emissions (Piva et al., 2014).

Overall, low N₂O fluxes in the eucalypt component could be a consequence of the high demand for nitrogen by the plants, reducing the loss of mineral nitrogen as N₂O, helping to mitigate the emissions (Figueiredo et al., 2018). Moreover, the low N₂O fluxes in these forest component systems may be related to changes in the phenolic content of eucalypt litter (Soumare et al., 2015), which contributes to the decreasing microbial community size and enzymatic activities and the increasing physiological microbial stress (Chen et al., 2013). Carvalho et al. (2017) evaluated ICLF and integrated crop–livestock (ICL) system over a 2-year period and found that soil N₂O fluxes were lower in the ICLF system than in the ICL system associated with eucalypt litter, which is rich in phenolic compounds that leads to a low carbon microbial biomass.

Materials and Methods

Study area characterization

The experimental field was located in the municipality of Sinop, Mato Grosso, Brazil (11°51 'S, 55°35 'W, 384 m altitude) (Figure 3) in the experimental field of Embrapa Agrossilvipastoril. The climate is classified as Am (Köppen monsoon climate) (Alvares et al., 2013). The annual average air temperature is 25.8 °C with an accumulated rainfall of 2,250 mm (Embrapa, 2017), with a dry season that extends from May to September (Souza et al., 2013). The average annual relative humidity is 71% (Embrapa, 2017).

The soil of the experimental area was classified as a typical dystrophic red-yellow latosol according to the Brazilian Soil Taxonomy (Santos et al., 2018), which is equivalent to a Hapludox under the US Soil Taxonomy (Soil Survey Staff, 2014). It has a clayey texture, moderately flat relief, subperennial vegetation, and kaolinitic-gibbsite composition (Viana et al., 2015). The soil, in the 0–20 cm layer, presented the following features at the start of the experiment: 5.7 pH in H₂O, 13.7 mg dm⁻³ of P (Mehlich 1), 79 mg dm⁻³ of K (Mehlich 1); 2.3 cmolc dm-3 of Ca, 0.66 cmolc dm⁻³ of Mg, 0.01 cmolc dm⁻³ of Al, and 29.6 g kg⁻¹ of organic matter (Farias Neto et al., 2019). The soil texture was clayey with 28% sand, 16% silt, and 56% clay. The experimental field originally showed a transition between the Cerrado and Amazon vegetation. Deforestation occurred in 1984 for the production of cassava (*Manihot esculenta* Crantz) (Araujo et al., 2009). In the 1990s, it was cultivated with rice (*Oryza sativa*) and later with soybean (*Glycine max* L.). Between 2002 and 2007, the area was cultivated with soybean and corn (*Zea mays* L.), and during the 2007–2009 harvests, soybean and cotton (*Gossypium hirsutum* L.) successions were conducted. During the 2010–2011 harvest, the area remained fallow (Diel et al., 2014).

In October 2011 (2011–2012 season), four treatments were implemented using a randomized complete block design with three replications. There were three types of integrated crop-livestock-forest (ICLF) systems: livestock-forest (LF), livestock-crop-forest (LCF), and ICLF, named in the order of entry of the agricultural component and livestock in each treatment, and a eucalypt monoculture (F). The F was evaluated in experimental plots of 1 ha and the other treatments (LF, LCF, and ICLF) were in plots of 2 ha, amounting to a total area of 21 ha with 12 experimental units (Figure 4).

The hybrid Eucalyptus urophylla × E. grandis clone H13 was used in the F treatment, and the clones were implanted with a spacing of 3.5×3.0 m and a density of 952 plants ha⁻¹. In the other treatments (LF, LCF, and ICLF), the clones were planted with a spacing of 3.5×3.0 m, in the east-west direction, and with a distance of 30 m between each grove. Furthermore, each grove consisted of three rows of eucalypt trees, totaling 270 plants ha⁻¹. LF treatment always consisted of Brachiaria brizantha (Hochst. ex A. Rich.) R. D. Webster grass between eucalypt groves. In the LCF treatment, crop and livestock components were alternated every 2 years. In the first two harvests (2011-2012 and 2012-2013), marandu palisade grass (Brachiaria brizantha 'Marandu') was cultivated between eucalypt groves. In the third and fourth harvest years (2013-2014 and 2014-2015), the area between groves was planted with soybean [Glycine max L. (Merr.)] from October to February, followed by corn (Zea mays L.) in combination with marandu palisade grass (cover crop) from February to July. In the ICLF treatment, the crop and livestock components were alternated annually. Soybean was cultivated every year between October and February (2011-2012, 2013-2014), followed by corn in combination with marandu palisade grass (cover crop) from February to July (Figure 5).

The planting of eucalypt in all treatments was carried out in November 2011. Plowing was performed at a depth of 50 cm, and 350 kg ha⁻¹ of simple superphosphate was applied to the planting furrow.

The activities of each treatment varied over the first three years of the system in relation to culture and fertilization, as shown in Figure 5.

The treatments were established in 2011, and until 2015, there were no livestock insertions. That is, during the period of evaluation of this research, no animal grazing occurred in any of the treatments, as the intention was to use the pasture for cutting in the use of silage and as a cover crop during the first year. As the experimental plot is located in an Amazon region with high rainfall, soil cover and enhancement of organic matter in the soil for other crops is extremely essential. Therefore, in the beginning of the experiment the animals were not inserted into the area, and the marandu palisade grass was used for silage and hay, being harvested with mechanical machines. Gas samples were collected every seven days during the 2013–2014 harvest (November 2013 to October 2014). Owing to logistics issues related to the circulation of agricultural machinery, it was not possible to perform sampling in February 2014. Thus, the fluxes from this month could not be measured.

 N_2O flux was collected through a manual static chamber in a base-top rectangular model, similar to that described by Nogueira et al. (2015). Metal bases (length × width × height) of 60 × 40 × 11 cm were installed in the field one week before the first collection in order to avoid disturbances in the soil. The bases were fixed at a depth of 5 cm in the soil and were maintained throughout the experiment. The top of the chamber consisted of a 60 × 40 × 9.2 cm (length × width × height) polyvinyl chloride (PVC) tray covered with a double-sided thermal blanket to decrease solar absorption and maintain internal temperature (Parkin and Venterea, 2010). Approximately 50 cm³ of air that had accumulated in the chamber was removed through a three-way valve connected to a 60 cm³ polypropylene syringe.

After chamber coupling (i.e., bottom-top), an air sample was immediately taken from the interior of the chamber, considered as time zero, and subsequent samples were taken at 20, 40, and 60 min (Parkin and Venterea, 2010). Consequently, the tops were removed from the base and the soil area was again exposed to the environmental conditions. N_2O sampling was always performed between 8 and 11 h (Zuchello, 2010). At the time of gas collection, the internal chamber temperature used for the N_2O flux calculation was monitored using thermohygrometers.

In each treatment, a chamber was allocated to the soil under the eucalypt tree canopy. In the F treatment, the chamber was installed at the center of it. The chambers were installed in the central row of the central eucalypt grove in the integrated systems (e.g., LF, LCF, and ICLF). The static chambers evaluated soils from within the eucalypt component of the respective systems, and not in areas with crop or livestock components. The meteorological station of Embrapa (Embrapa, 2017) was used to obtain the daily accumulated rainfall (mm), and then the accumulated monthly rainfall was calculated.

Laboratory analysis

Gases were transferred from syringes to butyl rubber sealed vials (e.g., glass vial) with 20 cm³ (Parkin and Venterea, 2010), previously vacuumed (approximately -100 kPa) using an electric vacuum pumping system for concentration analysis. After the transfer of the gases, the vials were left in an air-conditioned room at a temperature of 21 °C for approximately 24 h before analysis.

Subsequently, the samples were analyzed using a gas chromatography (Shimadzu GC-2014) equipped with an automatic injector and electron capture detector (ECD) to measure N₂O concentrations with 95% precision. The ECD detector temperature was maintained at 325 °C, and the column was maintained at 75 °C in the isothermal system. The columns were in the Hayesep series (1.0, 4.0 M; 1.5 M; 1.5 M; 0.7 M). Ultrapure nitrogen was used as the carrier gas at a flux rate of 25 mL min⁻¹, and the injector pressure was maintained at 300 kPa. To determine the standard curve, three gaseous solutions containing the N₂O standard (white Martins) were used. The default values were 382.8, 808, and 2080 mol mol⁻¹, respectively. The sample run time was eight minutes.

From the sample concentrations, the gas concentration change rate was calculated by considering the linear fit model. After obtaining the best fit in the gas increment inside the chamber, the flux was determined according to Equation 1, proposed by Hutchinson and Livingston (Hutchinson and Livingston, 1993). From equation 1, the fluxes of N₂O in μ g N m⁻² h⁻¹ were obtained, and the cumulative emission (kg N-N₂O ha⁻¹) was obtained.

$N_2 OF lows(\mu g N m^2 h^{-1}) = \frac{\left(\frac{\Delta c}{\Delta t}\right)^{*V}}{A} * \left(\frac{m}{Vm}\right)$ where $\Delta C/\Delta t$ is the slope of a linear function adjusted to the

where $\Delta C/\Delta t$ is the slope of a linear function adjusted to the gas concentration of the samples taken at 0, 20, 40, and 60 min after chamber closure (ppm hour⁻¹); V is the chamber volume (L), A is the chamber area (m²), m is the gas molecular weight (g mol⁻¹), and Vm is the molar volume of the gas (m³ mol⁻¹) corrected for air temperature (K) from inside the chamber.

Data analysis

The cumulative N_2O emissions (kg N-N₂O ha⁻¹) of dry (May to September) and wet (October to April) seasons were estimated using the trapezoidal integration method (Hergoualc'h et al., 2019). The net cumulative emissions from the entire period were estimated by subtracting the wet and dry period emissions. The data were not normally distributed, and nonparametric statistics were used. We used the Kruskal–Wallis (Kruskal and Wallis, 1952) test to perform comparisons among treatments by season, and Mann–Whitney U-test (Mann and Whitney, 1947) to perform comparisons between dry and wet seasons. All data analyses were performed using R Core Software (R Core Team, 2016).

Conclusions

Soil use and management intensification in other components from the ICLF system did not increase N_2O emission from the eucalypt soil, which was equal to that of eucalypt in monoculture (in which the total area is less intensive). It was observed that across all the treatments, eucalypt soil had positive emission values in the wet season, whereas in the dry season, the soil acted as a sink with negative emission values, thereby proving the influence of rainfall on the N_2O emissions. ICLF is considered a technological option that contributes to the objectives of the ABC Plan to Brazil to achieve its NDCs commitment under the Paris Agreement. However, to confirm the mitigation potential of ICLF systems, it is necessary to quantify N_2O emissions from all components of the system (crop, livestock, and forests).

Acknowledgments

We are grateful for the funding from the National Council for Scientific and Technological Development (CNPq) (grant number 472653/2013-0) and the Climate Network. We also thank Embrapa Agrossilvipastoril for the support on the conduction of the work and Renato Marques for his support in the elaboration of the study area location figure. We also thanks the Editage (www.editage.com) for the language editing support.

References

Alvares CA, Stape JL, Sentelhas PC, de Moraes Gonçalves JL, Sparovek G (2013) Köppen's climate classification map for Brazil. Meteorol Z. 711–728.

- Alves BJ, Madari BE, Boddey RM (2017) Integrated crop–livestock– forestry systems: prospects for a sustainable agricultural intensification. Nutr Cycl Agroecosys. 108: 1–4.
- Amadi CC, Van Rees KCJ, Farrell RE (2016) Soil–atmosphere exchange of carbon dioxide, methane and nitrous oxide in shelterbelts compared with adjacent cropped fields. Agr Ecosyst Environ. 223: 123–134.
- Araujo R de A, Costa RB da, Felfili JM, Gonçalvez IK, Sousa RAT de M, Dorval A (2009) Florística e estrutura de fragmento florestal em área de transição na Amazônia Matogrossense no município de Sinop. Acta Amazon. 39: 865–877.
- Bateman EJ, Baggs EM (2005) Contributions of nitrification and denitrification to N2O emissions from soils at different water-filled pore space. Biol Fert Soils. 41: 379–388.
- Bauters M, Verbeeck H, Rütting T, Barthel M, Mujinya BB, Bamba F, Bodé S, Boyemba F, Bulonza E, Carlsson E, Eriksson L, Makelele I, Six J, Ntaboba LC, Boeckx P (2019) Contrasting nitrogen fluxes in African tropical forests of the Congo Basin. Ecol Monogr. 89: e01342.
- Bayer C, Gomes J, Zanatta JA, Vieira FCB, Piccolo M de C, Dieckow J, Six J (2015) Soil nitrous oxide emissions as affected by long-term tillage, cropping systems and nitrogen fertilization in Southern Brazil. Soil Till Res. 146: 213–222.
- Benoit M, Garnier J, Billen G, Tournebize J, Gréhan E, Mary B (2015) Nitrous oxide emissions and nitrate leaching in an organic and a conventional cropping system (Seine basin, France). Agr Ecosyst Environ. 213: 131–141.
- Bouwman AF (1998) Nitrogen oxides and tropical agriculture. Nature. 392: 866–867.
- Brasil (2015) Brazilian Intended Nationally Determined Contribution to achieving the objective of the United Nations Framework Convention on Climate Change. Ministry of Foreign Affairs, Brasília.
- Brasil (2015). Ministério da Ciência, Tecnologia, Inovações e Comunicações (MCTI), Secretaria de Políticas para a Formação e Ações Estratégicas. Estimativas anuais de emissões de gases de efeito estufa no Brasil, Coordenação-Geral do Clima, 5^{rd.} edn. Brasília: Ministério da Ciência, Tecnologia, Inovações e Comunicações. 71.
- Burrows WH, Henry BK, Back PV, Hoffmann MB, Tait LJ, Anderson ER, Menke N, Danaher T, Carter JO, McKeon GM (2002) Growth and carbon stock change in eucalypt woodlands in northeast Australia: ecological and greenhouse sink implications. Glob Change Biol. 8: 769–784.
- Butterbach-Bahl K, Baggs EM, Dannenmann M, Kiese R, Zechmeister-Boltenstern S (2013) Nitrous oxide emissions from soils: how well do we understand the processes and their controls? Phil Trans R Soc B. 368: 20130122.
- Carvalho AM de, Oliveira WRD de, Ramos MLG, Coser TR, Oliveira AD de, Pulrolnik K, Souza KW, Vilela L, Marchão RL (2017) Soil N₂O fluxes in integrated production systems, continuous pasture and Cerrado. Nutr Cycl Agroecosys. 108: 69–83.
- Carvalho JLN, Raucci GS, Frazão LA, Cerri CEP, Bernoux M, Cerri CC (2014) Crop-pasture rotation: A strategy to reduce soil greenhouse gas emissions in the Brazilian Cerrado.Agr Ecosyst Environ. 183: 167–175.
- Cerri CEP, Easter M, Paustian K, Killian K, Coleman K, Bernoux M, Falloon P, Powlson DS, Batjes N, Milne E, Cerri CC (2007) Simulating SOC changes in 11 land use change chronosequences from the Brazilian Amazon with RothC and Century models. Agr Ecosyst Environ. 122: 46–57.
- Chapuis-Lardy L, Wrage N, Metay A, Chotte J-L, Bernoux M (2007) Soils, a sink for N₂O? A review. Glob Chang Biol. 13: 1–17.
- Charteris AF, Chadwick DR, Thorman RE, Vallejo A, De Klein CAM, Rochette P, Cárdenas LM (2020) Global Research Alliance N_2O Chamber Methodology Guidelines: Recommendations for deployment and accounting for sources of variability. J Environ Qual. 49:1092–1109.
- Chen F, Zheng H, Zhang K, Ouyang Z, Li H, Wu B, Shi Q (2013) Soil microbial community structure and function responses to successive planting of Eucalyptus. J Environ Sci. 25: 2102–2111.
- Cordeiro LAM, Vilela L, Marchão RL, Kluthcouski J, Júnior GBM (2015) Integração lavoura-pecuária e integração lavoura-pecuária-

floresta: estratégias para intensificação sustentável do uso do solo. CC&T. 32: 15–53.

- Corrêa RS, Madari BE, Carvalho GD, Costa AR da, Pereira AC de C, Medeiros JC (2016) Fluxos de óxido nitroso e suas relações com atributos físicos e químicos do solo. Pesqui Agropecu Bras. 51: 1148–1155.
- Cuer CA, Rodrigues R de AR, Balieiro FC, Jesus J, Silva EP, Alves BJR, Rachid CTCC (2018) Short-term effect of Eucalyptus plantations on soil microbial communities and soil-atmosphere methane and nitrous oxide exchange. Sci Rep-UK. 8: 1–12.
- Davidson EA, Keller M, Erickson HE, Verchot LV, Veldkamp E (2000) Testing a Conceptual Model of Soil Emissions of Nitrous and Nitric Oxides. BioScience. 50: 667–680.
- Diel D, Behling M, Neto AL de F, Isernhagen ECC (2014) Distribuição horizontal e vertical de fósforo em sistemas de cultivos exclusivos de soja e de integração lavoura-pecuária-floresta. Pesqui Agropecu Bras. 49: 639–647.
- Du H, Zeng F, Peng W, Wang K, Zhang H, Liu L, Song T (2015) Carbon Storage in a Eucalyptus Plantation Chronosequence in Southern China. Forests. 6: 1763–1778.
- Dube F, Espinosa M, Stolpe NB, Zagal E, Thevathasan NV, Gordon AM (2012) Productivity and carbon storage in silvopastoral systems with Pinus ponderosa and Trifolium spp., plantations and pasture on an Andisol in Patagonia, Chile. Agroforest Syst. 86: 113–128.
- Embrapa (2017) Estação Meteorológica Portal Embrapa. Available in: https://www.embrapa.br/agrossilvipastoril/estacaometeorologica>. Accessed 3rd June, 2018.
- Euclides VPB, do Valle CB, Macedo MCM, Almeida RG de, Montagner DB, Barbosa RA (2010) Brazilian scientific progress in pasture research during the first decade of XXI century. Rev Bras Zootecn. 39: 151–168.
- Fang L, Cong-Qiang L, Shi-lu W, Zheng-jie Z (2012) Soil temperature and moisture controls on surface fluxes and profile concentrations of greenhouse gases in karst area in central part of Guizhou Province, southwest China. Environ Earth Sci. 67: 1431– 1439.
- Farias Neto AL, Silva JFV, Meneguci JLP, Magalhães CAS, Pedreira BC, Tonini H, Matos ES, Behling M, Ferreira A (2019) Estabelecimento de Sistemas Integração Lavoura- Pecuária-Floresta com foco em gado de corte na Embrapa Agrossilvipastoril. In: Farias Neto AL de, Nascimento AF do, Rossoni AL, Magalhães CA de S, Ituassu DR, Hoogerheide ESS, Ikeda FS, Fernandes Junior F, Faria GR, Isernhagen I, Vendrusculo LG, Morales MM, Carnevalli RA Embrapa Agrossilvipastoril: primeiras contribuições para o desenvolvimento de uma agropecuária sustentável, 1rdedn. Brasília: Embrapa. 145-163.
- Figueiredo EB, Jayasundara S, de Oliveira Bordonal R, Berchielli TT, Reis RA, Wagner-Riddle C, La Scala Jr. N (2017) Greenhouse gas balance and carbon footprint of beef cattle in three contrasting pasture-management systems in Brazil. J Clean Prod. 142: 420– 431.
- Figueiredo CC, de Oliveira AD, dos Santos IL, Ferreira EAB, Malaquias JV, de Sá MAC, de Carvalho AM, dos Santos J de DG (2018) Relationships between soil organic matter pools and nitrous oxide emissions of agroecosystems in the Brazilian Cerrado. Sci Total Environ. 618: 1572–1582.
- Gil J, Siebold M, Berger T (2015) Adoption and development of integrated crop–livestock–forestry systems in Mato Grosso, Brazil. Agr Ecosyst Environ. 199: 394–406.
- Hartmann DL, Klein Tank AMG, Rusticucci M, Alexander LV, Brönnimann S, Charabi Y, Dentener FJ, Dlugokencky EJ, Easterling DR, Kaplan A, Soden BJ, Thorne PW, Wild M, Zhai PM (2013) Observations: Atmosphere and Surface. In: Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 159–254.
- Hergoualc'h K, Akiyama H, Bernoux M, Chirinda N, Prado A del, Kasimir Å, MacDonald JD, Ogle SM, Regina K, van der Weerden TJ, Liang C, Noble A, (2019) N₂O Emissions From Managed Soils, and CO₂ Emissions From Lime and Urea Application. In: Calvo Buendia

E, Tanabe K, Kranjc A, Baasansuren J, Fukuda M, Ngarize S, Osako A, Pyrozhenko Y, Shermanau P, Federici S. (Eds.) Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. IPCC, Switzerland. 1–48.

- Hutchinson GL, Livingston GP (1993) Use of Chamber Systems to Measure Trace Gas Fluxes. In: Rolston DE, Duxbury JM, Harper LA, Mosier AR (eds.) Agricultural Ecosystem Effects on Trace Gases and Global Climate Change, ASA Special Publications, 55th edn. American Society of Agronomy, Crop Science Society of America and Soil Science Society of America, Madison, USA. 63-78
- Jain N, Arora P, Tomer R, Mishra SV, Bhatia A, Pathak H, Chakraborty D, Kumar V, Dubey DS, Harit RC, Singh JP (2016) Greenhouse gases emission from soils under major crops in Northwest India. Sci Total Environ. 542: 551–561.
- Konda R, Ohta S, Ishizuka S, Arai S, Ansori S, Tanaka N, Hardjono A (2008) Spatial structures of N2O, CO2, and CH4 fluxes from Acacia mangium plantation soils during a relatively dry season in Indonesia. Soil Biol Biochem. 40: 3021–3030.
- Kruskal WH, Wallis WA (1952) Use of Ranks in One-Criterion Variance Analysis. J Am Stat Assoc. 47: 583–621.
- Li S, Gu X, Zhuang J, An T, Pei J, Xie H, Li H, Fu S, Wang J (2016) Distribution and storage of crop residue carbon in aggregates and its contribution to organic carbon of soil with low fertility. Soil Till Res. 155: 199–206.
- Linn DM, Doran JW (1984) Effect of Water-Filled Pore Space on Carbon Dioxide and Nitrous Oxide Production in Tilled and Nontilled Soils. Soil Sci Soc Am J. 48: 1267–1272.
- Mann HB, Whitney DR (1947) On a Test of Whether one of Two Random Variables is Stochastically Larger than the Other. The Annals of Mathematical Statistics, Institute of Mathematical Statistics. 18: 50–60.
- Martins MR, Jantalia CP, Polidoro JC, Batista JN, Alves BJR, Boddey RM, Urquiaga S (2015) Nitrous oxide and ammonia emissions from N fertilization of maize crop under no-till in a Cerrado soil. Soil Till Res. 151: 75–81.
- McDaniel MD, Simpson RR, Malone BP, McBratney AB, Minasny B, Adams MA (2017) Quantifying and predicting spatio-temporal variability of soil CH4 and N_2O fluxes from a seemingly homogeneous Australian agricultural field. Agr Ecosyst Environ. 240:182–193.
- Muller F (2005) Indicating ecosystem and landscape organisation. Ecol Indic. 5: 280–294.
- Myhre G, Shindell D, Bréon F-M, Collins W, Fuglestvedt J, Huang J, Koch D, Lamarque J-F, Lee D, Mendoza B, Nakajima T, Robock A, Stephens G, Takemura T, Zhang H (2013) Anthropogenic and Natural Radiative Forcing. In: Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. 659–740.
- Nair PKR (2012) Climate Change Mitigation: A Low-Hanging Fruit of Agroforestry. In: Nair PKR, Garrity D (eds) Agroforestry - The Future of Global Land Use, Springer Netherlands: Dordrecht. 31– 67.
- Nascimento AF do, Rodrigues R de AR (2019) Sampling frequency to estimate cumulative nitrous oxide emissions from the soil. Pesqui Agropecu Bras. 54: e00211.
- Nascimento AF do, Rodrigues R de AR, Silveira JG da, Silva JJN da, Daniel V de C, Segatto ER (2020) Nitrous oxide emissions from a tropical Oxisol under monocultures and an integrated system in the Southern Amazon – Brazil. Rev Bras Cienc Solo. 44: e0190123.
- Neves APR (2016) Dinâmica de gases de efeito estufa em plantações de *Acacia mearnsii* De Wildem solos no Bioma Pampa. Mestrado, Universidade Federal de Pelotas, Rio Grande do Sul.
- Nogueira AK da S, Rodrigues R de AR, Castro BS, Nogueira TF, Silva JJN da, Behling M, Mombach M, Armacolo N, Silveira JG (2015) Emissões de óxido nitroso e metano do solo em áreas de recuperação de pastagens na Amazônia matogrossense. Quim Nova. 38: 937–943.
- Nogueira AK da S, Rodrigues R de AR, Silva JJN da, Botin AA, Silveira JG da, Mombach MA, Armacolo NM, Romeiro S de O (2016) Fluxos de óxido nitroso em sistema de integração lavoura-

pecuária-floresta. Pesqui Agropecu Bras. 51: 1156–1162.

- Parkin T, Venterea R (2010) USDA-ARS GRACEnet Project Protocols Chapter 3. Chamber-Based Trace Gas Flux Measurements. Sampling Protocols.
- Paula RR, Reis GG, Reis MGF, Oliveira Neto SN, Leite HG, Melido RCN, Lopes HNS, Souza FC (2013) Eucalypt growth in monoculture and silvopastoral systems with varied tree initial densities and spatial arrangements. Agroforest Syst. 87: 1295–1307.
- Paustian K, Lehmann J, Ogle S, Reay D, Robertson GP, Smith P (2016) Climate-smart soils. Nature. 532: 49–57.
- Piva JT, Dieckow J, Bayer C, Zanatta JA, de Moraes A, Tomazi M, Pauletti V, Barth G, Piccolo M de C (2014) Soil gaseous N_2O and CH4 emissions and carbon pool due to integrated crop-livestock in a subtropical Ferralsol. Agr Ecosyst Environ. 190: 87–93.
- Piva JT, Sartor LR, Sandini IE, Moraes A de, Dieckow J, Bayer C, Rosa CM da (2019) Emissions of Nitrous Oxide and Methane in a Subtropical Ferralsol Subjected to Nitrogen Fertilization and Sheep Grazing in Integrated Crop-Livestock System. Rev Bras Cienc Solo. 43: e0180140.
- R Core Team (2016) 'R: A Language and Environment for Statistical Computing.' (R Foundation for Statistical Computing: Vienna, Austria). Available in: https://www.R-project.org/. Accessed 5rd May, 2020.
- Reay DS, Davidson EA, Smith KA, Smith P, Melillo JM, Dentener F, Crutzen PJ (2012) Global agriculture and nitrous oxide emissions. Nat Clim Change. 2: 410–416.
- Rede ILPF (2016) ILPF em Números Portal Embrapa. Available in: https://www.embrapa.br/web/rede-ilpf/ilpf-em-numeros. Accessed 5rd May, 2020.
- Rivera JE, Chará J, Barahona R (2019) CH4, CO2 and N₂O emissions from grasslands and bovine excreta in two intensive tropical dairy production systems. Agroforest Syst. 93:915–928.
- Rodrigues R de AR, Mello WZ de, Conceição MCG da, Souza PA de, Silva JJN da (2017) Dinâmica do nitrogênio em sistemas agrícolas e florestais tropicais e seu impacto na mudança do clima. Rev Virtual Quím. 9: 1868–1886.
- Rosenkranz P, Brüggemann N, Papen H, Xu Z, Horváth L, Butterbach-Bahl K (2006) Soil N and C trace gas fluxes and microbial soil N turnover in a sessile oak (Quercus petraea (Matt.) Liebl.) forest in Hungary. Plant Soil. 286: 301–322.
- Saggar S (2010) Estimation of nitrous oxide emission from ecosystems and its mitigation technologies. Agr Ecosyst Environ. 136: 189–191.
- Salton JC, Mercante FM, Tomazi M, Zanatta JA, Concenço G, Silva WM, Retore M (2014) Integrated crop-livestock system in tropical Brazil: Toward a sustainable production system. Agr Ecosyst Environ. 190: 70–79.
- Santos HG dos, Jacomine PKT, Anjos LHC dos, Oliveira VÁ de, Lumbreras JF, Coelho MR, Almeida JA de, Filho JC de A, Oliveira JB de, Cunha TJF (2018) Sistema Brasileiro de Classificação de Solos, 5rd edn. Brasília, DF: Embrapa.
- Sato JH, de Carvalho AM, de Figueiredo CC, Coser TR, de Sousa TR, Vilela L, Marchão RL (2017) Nitrous oxide fluxes in a Brazilian clayey oxisol after 24 years of integrated crop-livestock management. Nutr Cycl Agroecosys. 108: 55–68.
- Scott CE, Monks SA, Spracklen DV, Arnold SR, Forster PM, Rap A, Äijälä M, Artaxo P, Carslaw KS, Chipperfield MP, Ehn M, Gilardoni S, Heikkinen L, Kulmala M, Petäjä T, Reddington CLS, Rizzo LV, Swietlicki E, Vignati E, Wilson C (2018) Impact on short-lived climate forcers increases projected warming due to deforestation. Nature. 9: 157.
- Silva JJN da (2019) Dinâmica do nitrogênio com enfoque nas emissões de N_2O em plantio comercial de Eucalipto no bioma da Mata Atlântica. Doutorado, Universidade Federal Fluminense, Rio de Janeiro.
- Silveira N, Piatto M, Costa C, Medeiros M, Pinto LF (2018) SEEG -Relatório Analítico Agropecuária. Available in: https://seeg.eco.br/>. Accessed 10rd July, 2020.
- Soil Survey Staff (2014) Keys to Soil Taxonomy | NRCS Soils. (USDA-Natural Resources Conservation Service: Washington, DC). Available in: https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/ class/taxonomy/?cid=nrcs142p2_053580>. Accessed 10rd July, 2020.

- Soumare A, Manga A, Fall S, Hafidi M, Ndoye I, Duponnois R (2015) Effect of Eucalyptus camaldulensis amendment on soil chemical properties, enzymatic activity, Acacia species growth and roots symbioses. Agroforest Syst. 89: 97–106.
- Souza AP, Mota LL, Zamadei T, Martim CC, Almeida FT, Paulino J (2013) Classificação Climática e Balanço Hídrico Climatológico no Estado de Mato Grosso. Nativa. 1: 34–43.
- Stocker TF, Qin D, Plattner GK, Alexander LV, Allen SK, Bindoff NL, Bréon FM, Church AJ, Cubasch U, Emori S, Forster P, Friedlingstein P, Gillett N, Gregory JM, Hartmann DL, Jansen E, Kirtman B, Knutti R, Krishna Kumar K, Lemke P, Marotzke J, Masson-Delmotte V, Meehl GA, Mokhov II, Piao S, Ramaswamy V, Randall D, Rhein M, Rojas M, Sabine C, Shindell D, Talley LD, Vaughan DG, Xie SP (2013) Technical Summary. In: Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [(eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Thevathasan NV, Gordon AM, Bradley R, Cogliastro A, Folkard P, Grant R, Kort J, Liggins L, Njenga F, Olivier A, Pharo C, Powell G, Rivest D, Schiks T, Trotter D, Van Rees K, Whalen J, Zabek L (2012) Agroforestry Research and Development in Canada: The Way Forward. Agroforestry - The Future of Global Land Use. (Eds PKR Nair, D Garrity). Adv Agrofor. pp. 247–283. (Springer Netherlands: Dordrecht).
- Tian H, Chen G, Lu C, Xu X, Ren W, Zhang B, Banger K, Tao B, Pan S, Liu M, Zhang C, Bruhwiler L, Wofsy S (2015) Global methane and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental changes. Ecosystem Health and Sustainability. 1: 1–20.
- Tian H, Xu R, Canadell JG, Thompson RL, Winiwarter W, Suntharalingam P, Davidson EA, Ciais P, Jackson RB, Janssens-Maenhout G, Prather MJ, Regnier P, Pan N, Pan S, Peters GP, Shi H, Tubiello FN, Zaehle S, Zhou F, Arneth A, Battaglia G, Berthet S, Bopp L, Bouwman AF, Buitenhuis ET, Chang J, Chipperfield MP, Dangal SRS, Dlugokencky E, Elkins JW, Eyre BD, Fu B, Hall B, Ito A, Joos F, Krummel PB, Landolfi A, Laruelle GG, Lauerwald R, Li W, Lienert S, Maavara T, MacLeod M, Millet DB, Olin S, Patra PK, Prinn RG, Raymond PA, Ruiz DJ, van der Werf GR, Vuichard N, Wang J, Weiss RF, Wells KC, Wilson C, Yang J, Yao Y (2020) A comprehensive quantification of global nitrous oxide sources and sinks. Nature. 586: 248–256.
- Ussiri D, Lal R (2013) Formation and Release of Nitrous Oxide from Terrestrial and Aquatic Ecosystems. In: Ussiri D, Lal R (eds) Soil Emission of Nitrous Oxide and its Mitigation, Springer Netherlands: Dordrecht. 63–96.
- Viana JHM, Spera ST, Magalhaes CA de S, Calderano SB (2015) Caracterização dos solos do sítio experimental dos ensaios do Projeto Safrinha em Sinop-MT. Comunicado Técnico EMBRAPA. 210.
- Weerden TJ van der, Kelliher FM, Klein CAM de (2012) Influence of pore size distribution and soil water content on nitrous oxide emissions. Soil Res. 50: 125–135.
- WMO Greenhouse Gas Bulletin (2018) The State of Greenhouse
 Gases in the Atmosphere Based on Global Observations through

 2017.
 Available
 in:
- <https://library.wmo.int/doc_num.php?explnum_id=5455>.
 Accessed 8rd August, 2020.
- Zhang K, Zheng H, Chen F, Li R, Yang M, Ouyang Z, Lan J, Xiang X (2017) Impact of nitrogen fertilization on soil–Atmosphere greenhouse gas exchanges in eucalypt plantations with different soil characteristics in southern China. Plos One. 12: 1-14.
- Zomer RJ, Neufeldt H, Xu J, Ahrends A, Bossio D, Trabucco A, Noordwijk M van, Wang M (2016) Global Tree Cover and Biomass Carbon on Agricultural Land: The contribution of agroforestry to global and national carbon budgets. Sci Rep-UK. 6: 1–12.
- Zuchello F (2010) Emissões de óxido nitroso (N2O) de Cambissolo cultivado com cana-de-açúcar em Campos dos Goytacazes: impacto de adubações com vinhaça e uréia. Doutorado, Universidade Federal Rural do Rio de Janeiro, Rio de Janeiro.